

**INVOKANA®**

(canagliflozin) tablets, for oral use

Revised: 07/2016  
058051-160811**HIGHLIGHTS OF PRESCRIBING INFORMATION**

These highlights do not include all the information needed to use INVOKANA® safely and effectively. See full prescribing information for INVOKANA.

INVOKANA (canagliflozin) tablets, for oral use  
Initial U.S. Approval: 2013

**RECENT MAJOR CHANGES**

Dosage and Administration (2.2)	05/2016
Contraindications (4)	03/2016
Warnings and Precautions (5)	07/2016

**INDICATIONS AND USAGE**

INVOKANA is a sodium-glucose co-transporter 2 (SGLT2) inhibitor indicated as an adjunct to diet and exercise to improve glycemic control in adults with type 2 diabetes mellitus (1)

**Limitation of Use:**

- Not for treatment of type 1 diabetes mellitus or diabetic ketoacidosis (1)

**DOSE AND ADMINISTRATION**

- The recommended starting dose is 100 mg once daily, taken before the first meal of the day (2.1)
- Dose can be increased to 300 mg once daily in patients tolerating INVOKANA 100 mg once daily who have an eGFR of 60 mL/min/1.73 m<sup>2</sup> or greater and require additional glycemic control (2.1)
- Assess renal function before initiating and periodically thereafter. (2.2)
- Limit the dose of INVOKANA to 100 mg once daily in patients who have an eGFR of 45 to less than 60 mL/min/1.73 m<sup>2</sup> (2.2)
- Initiation or use of INVOKANA is not recommended if eGFR is below 45 mL/min/1.73 m<sup>2</sup> (2.2)

**DOSE FORMS AND STRENGTHS**

Tablets: 100 mg, 300 mg (3)

**CONTRAINDICATIONS**

- History of serious hypersensitivity reaction to INVOKANA (4)
- Severe renal impairment, ESRD, or on dialysis (4)

**WARNINGS AND PRECAUTIONS**

- **Hypotension:** Before initiating INVOKANA, assess volume status and correct hypovolemia in patients with renal impairment, the elderly, in patients with low systolic blood pressure, or if on diuretics, ACEI, or ARB. Monitor for signs and symptoms during therapy (5.1)

**INVOKANA® (canagliflozin) tablets**

- **Ketoacidosis:** Assess patients who present with signs and symptoms of metabolic acidosis for ketoacidosis, regardless of blood glucose level. If suspected, discontinue INVOKANA, evaluate and treat promptly. Before initiating INVOKANA, consider risk factors for ketoacidosis. Patients on INVOKANA may require monitoring and temporary discontinuation of therapy in clinical situations known to predispose to ketoacidosis (5.2)
- **Acute kidney injury and impairment in renal function:** Consider temporarily discontinuing in settings of reduced oral intake or fluid losses. If acute kidney injury occurs, discontinue and promptly treat. Monitor renal function during therapy (5.3)
- **Hyperkalemia:** Monitor potassium levels in patients with impaired renal function and in patients predisposed to hyperkalemia (2.2, 5.4, 6.1, 8.6)
- **Urosepsis and Pyelonephritis:** Evaluate patients for signs and symptoms of urinary tract infections and treat promptly, if indicated (5.5)
- **Hypoglycemia:** Consider a lower dose of insulin or the insulin secretagogue to reduce the risk of hypoglycemia when used in combination with INVOKANA (5.6)
- **Genital mycotic infections:** Monitor and treat if indicated (5.7)
- **Hypersensitivity reactions:** Discontinue INVOKANA and monitor until signs and symptoms resolve (5.8)
- **Bone fracture:** Consider factors that contribute to fracture risk before initiating INVOKANA (5.9)
- **Increased LDL-C:** Monitor LDL-C and treat if appropriate (5.10)

**ADVERSE REACTIONS**

- Most common adverse reactions associated with INVOKANA (5% or greater incidence): female genital mycotic infections, urinary tract infection, and increased urination (6.1)

To report SUSPECTED ADVERSE REACTIONS, contact Janssen Pharmaceuticals, Inc. at 1-800-526-7738 or FDA at 1-800-FDA-1088 or [www.fda.gov/medwatch](http://www.fda.gov/medwatch).

**DRUG INTERACTIONS**

- **UGT Inducers** (e.g., rifampin): Canagliflozin exposure is reduced. Consider increasing dose from 100 mg to 300 mg (2.3, 7.1)
- **Digoxin:** Monitor digoxin levels (7.2)

**USE IN SPECIFIC POPULATIONS**

- **Pregnancy:** Advise females of the potential risk to a fetus especially during the second and third trimesters. (8.1)
- **Lactation:** INVOKANA is not recommended when breastfeeding (8.2)
- **Geriatrics:** Higher incidence of adverse reactions related to reduced intravascular volume (5.1, 8.5)
- **Renal Impairment:** Higher incidence of adverse reactions related to reduced intravascular volume and renal function (2.2, 5.3, 8.6)
- **Hepatic impairment:** Not recommended with severe hepatic impairment (8.7)

See 17 for PATIENT COUNSELING INFORMATION and Medication Guide.

Revised: 07/2016

**FULL PRESCRIBING INFORMATION: CONTENTS\***

- 1 INDICATIONS AND USAGE
- 2 DOSAGE AND ADMINISTRATION
  - 2.1 Recommended Dosage
  - 2.2 Patients with Renal Impairment
  - 2.3 Concomitant Use with UDP-Glucuronosyl Transferase (UGT) Enzyme Inducers
- 3 DOSAGE FORMS AND STRENGTHS
- 4 CONTRAINDICATIONS
- 5 WARNINGS AND PRECAUTIONS
  - 5.1 Hypotension
  - 5.2 Ketoacidosis
  - 5.3 Acute Kidney Injury and Impairment in Renal Function
  - 5.4 Hyperkalemia
  - 5.5 Urosepsis and Pyelonephritis
  - 5.6 Hypoglycemia with Concomitant Use with Insulin and Insulin Secretagogues
  - 5.7 Genital Mycotic Infections
  - 5.8 Hypersensitivity Reactions
  - 5.9 Bone Fracture
  - 5.10 Increases in Low-Density Lipoprotein (LDL-C)
  - 5.11 Macrovascular Outcomes
- 6 ADVERSE REACTIONS
  - 6.1 Clinical Studies Experience
  - 6.2 Postmarketing Experience

- 7 DRUG INTERACTIONS
  - 7.1 UGT Enzyme Inducers
  - 7.2 Digoxin
  - 7.3 Positive Urine Glucose Test
  - 7.4 Interference with 1,5-anhydroglucitol (1,5-AG) Assay
- 8 USE IN SPECIFIC POPULATIONS
  - 8.1 Pregnancy
  - 8.2 Lactation
  - 8.4 Pediatric Use
  - 8.5 Geriatric Use
  - 8.6 Renal Impairment
  - 8.7 Hepatic Impairment
- 10 OVERDOSAGE
- 11 DESCRIPTION
- 12 CLINICAL PHARMACOLOGY
  - 12.1 Mechanism of Action
  - 12.2 Pharmacodynamics
  - 12.3 Pharmacokinetics
- 13 NONCLINICAL TOXICOLOGY
  - 13.1 Carcinogenesis, Mutagenesis, Impairment of Fertility
- 14 CLINICAL STUDIES
  - 14.1 Monotherapy
  - 14.2 Combination Therapy
  - 14.3 Studies in Special Populations
- 16 HOW SUPPLIED/STORAGE AND HANDLING
- 17 PATIENT COUNSELING INFORMATION

\*Sections or subsections omitted from the full prescribing information are not listed.

**INVOKANA® (canagliflozin) tablets****FULL PRESCRIBING INFORMATION****1 INDICATIONS AND USAGE**

INVOKANA® (canagliflozin) is indicated as an adjunct to diet and exercise to improve glycemic control in adults with type 2 diabetes mellitus [see *Clinical Studies* (14)].

**Limitation of Use**

INVOKANA is not recommended in patients with type 1 diabetes mellitus or for the treatment of diabetic ketoacidosis.

**2 DOSAGE AND ADMINISTRATION****2.1 Recommended Dosage**

The recommended starting dose of INVOKANA (canagliflozin) is 100 mg once daily, taken before the first meal of the day. In patients tolerating INVOKANA 100 mg once daily who have an eGFR of 60 mL/min/1.73 m<sup>2</sup> or greater and require additional glycemic control, the dose can be increased to 300 mg once daily [see *Warnings and Precautions* (5.3), *Clinical Pharmacology* (12.2), and *Patient Counseling Information* (17)].

In patients with volume depletion, correcting this condition prior to initiation of INVOKANA is recommended [see *Warnings and Precautions* (5.1), *Use in Specific Populations* (8.5 and 8.6), and *Patient Counseling Information* (17)].

**2.2 Patients with Renal Impairment**

Assessment of renal function is recommended prior to initiation of INVOKANA and periodically thereafter.

The dose of INVOKANA is limited to 100 mg once daily in patients with moderate renal impairment with an eGFR of 45 to less than 60 mL/min/1.73 m<sup>2</sup>.

Initiation of INVOKANA is not recommended in patients with an eGFR less than 45 mL/min/1.73 m<sup>2</sup>.

Use of INVOKANA is not recommended when eGFR is persistently less than 45 mL/min/1.73 m<sup>2</sup> [see *Warnings and Precautions* (5.3) and *Use in Specific Populations* (8.6)].

INVOKANA is contraindicated in patients with an eGFR less than 30 mL/min/1.73 m<sup>2</sup> [see *Contraindications* (4)].

**2.3 Concomitant Use with UDP-Glucuronosyl Transferase (UGT) Enzyme Inducers**

If an inducer of UGTs (e.g., rifampin, phenytoin, phenobarbital, ritonavir) is co-administered with INVOKANA, consider increasing the dosage to 300 mg once daily in patients currently tolerating INVOKANA 100 mg once daily who have an eGFR of 60 mL/min/1.73 m<sup>2</sup> or greater and require additional glycemic control [see *Drug Interactions* (7.1)].

Consider another antihyperglycemic agent in patients with an eGFR of 45 to less than 60 mL/min/1.73 m<sup>2</sup> receiving concurrent therapy with a UGT inducer.

**3 DOSAGE FORMS AND STRENGTHS**

- INVOKANA 100 mg tablets are yellow, capsule-shaped, film-coated tablets with "CFZ" on one side and "100" on the other side.
- INVOKANA 300 mg tablets are white, capsule-shaped, film-coated tablets with "CFZ" on one side and "300" on the other side.

**4 CONTRAINDICATIONS**

- History of a serious hypersensitivity reaction to INVOKANA, such as anaphylaxis or angioedema [see *Warnings and Precautions* (5.8) and *Adverse Reactions* (6.1, 6.2)].
- Severe renal impairment (eGFR less than 30 mL/min/1.73 m<sup>2</sup>), end stage renal disease (ESRD), or patients on dialysis [see *Warnings and Precautions* (5.3) and *Use in Specific Populations* (8.6)].

**5 WARNINGS AND PRECAUTIONS****5.1 Hypotension**

INVOKANA causes intravascular volume contraction. Symptomatic hypotension can occur after initiating INVOKANA [see *Adverse Reactions* (6.1)] particularly in patients with impaired renal function (eGFR less than 60 mL/min/1.73 m<sup>2</sup>), elderly patients, patients on either diuretics or medications that interfere with the renin-angiotensin-aldosterone system (e.g., angiotensin-converting-enzyme [ACE] inhibitors, angiotensin receptor blockers [ARBs]), or patients with low systolic blood pressure. Before initiating INVOKANA in patients with one or more of these characteristics, volume status should be assessed and corrected. Monitor for signs and symptoms after initiating therapy.

**5.2 Ketoacidosis**

Reports of ketoacidosis, a serious life-threatening condition requiring urgent hospitalization have been identified in postmarketing surveillance in patients with type 1 and type 2 diabetes mellitus receiving sodium glucose co-transporter-2 (SGLT2) inhibitors, including INVOKANA. Fatal cases of ketoacidosis have been reported in patients taking INVOKANA. INVOKANA is not indicated for the treatment of patients with type 1 diabetes mellitus [see *Indications and Usage* (1)].

Patients treated with INVOKANA who present with signs and symptoms consistent with severe metabolic acidosis should be assessed for

**INVOKANA® (canagliflozin) tablets**

ketoacidosis regardless of presenting blood glucose levels, as ketoacidosis associated with INVOKANA may be present even if blood glucose levels are less than 250 mg/dL. If ketoacidosis is suspected, INVOKANA should be discontinued, patient should be evaluated, and prompt treatment should be instituted. Treatment of ketoacidosis may require insulin, fluid and carbohydrate replacement.

In many of the postmarketing reports, and particularly in patients with type 1 diabetes, the presence of ketoacidosis was not immediately recognized and institution of treatment was delayed because presenting blood glucose levels were below those typically expected for diabetic ketoacidosis (often less than 250 mg/dL). Signs and symptoms at presentation were consistent with dehydration and severe metabolic acidosis and included nausea, vomiting, abdominal pain, generalized malaise, and shortness of breath. In some but not all cases, factors predisposing to ketoacidosis such as insulin dose reduction, acute febrile illness, reduced caloric intake due to illness or surgery, pancreatic disorders suggesting insulin deficiency (e.g., type 1 diabetes, history of pancreatitis or pancreatic surgery), and alcohol abuse were identified.

Before initiating INVOKANA, consider factors in the patient history that may predispose to ketoacidosis including pancreatic insulin deficiency from any cause, caloric restriction, and alcohol abuse. In patients treated with INVOKANA consider monitoring for ketoacidosis and temporarily discontinuing INVOKANA in clinical situations known to predispose to ketoacidosis (e.g., prolonged fasting due to acute illness or surgery).

**5.3 Acute Kidney Injury and Impairment in Renal Function**

INVOKANA causes intravascular volume contraction [see *Warnings and Precautions* (5.1)] and can cause renal impairment [see *Adverse Reactions* (6.1)]. There have been postmarketing reports of acute kidney injury, some requiring hospitalization and dialysis, in patients receiving INVOKANA; some reports involved patients younger than 65 years of age.

Before initiating INVOKANA, consider factors that may predispose patients to acute kidney injury including hypovolemia, chronic renal insufficiency, congestive heart failure and concomitant medications (diuretics, ACE inhibitors, ARBs, NSAIDs). Consider temporarily discontinuing INVOKANA in any setting of reduced oral intake (such as acute illness or fasting) or fluid losses (such as gastrointestinal illness or excessive heat exposure); monitor patients for signs and symptoms of acute kidney injury. If acute kidney injury occurs, discontinue INVOKANA promptly and institute treatment.

INVOKANA increases serum creatinine and decreases eGFR. Patients with hypovolemia may be more susceptible to these changes. Renal function abnormalities can occur after initiating INVOKANA [see *Adverse Reactions* (6.1)]. Renal function should be evaluated prior to initiation of INVOKANA and monitored periodically thereafter. Dosage adjustment and more frequent renal function monitoring are recommended in patients with an eGFR below 60 mL/min/1.73 m<sup>2</sup>. Use of INVOKANA is not recommended when eGFR is persistently less than 45 mL/min/1.73 m<sup>2</sup> and is contraindicated in patients with an eGFR less than 30 mL/min/1.73 m<sup>2</sup> [see *Dosage and Administration* (2.2), *Contraindications* (4), *Use in Specific Populations* (8.6)].

**5.4 Hyperkalemia**

INVOKANA can lead to hyperkalemia. Patients with moderate renal impairment who are taking medications that interfere with potassium excretion, such as potassium-sparing diuretics, or medications that interfere with the renin-angiotensin-aldosterone system are at an increased risk of developing hyperkalemia [see *Dosage and Administration* (2.2) and *Adverse Reactions* (6.1)].

Monitor serum potassium levels periodically after initiating INVOKANA in patients with impaired renal function and in patients predisposed to hyperkalemia due to medications or other medical conditions.

**5.5 Urrosepsis and Pyelonephritis**

There have been postmarketing reports of serious urinary tract infections including urrosepsis and pyelonephritis requiring hospitalization in patients receiving SGLT2 inhibitors, including INVOKANA. Treatment with SGLT2 inhibitors increases the risk for urinary tract infections. Evaluate patients for signs and symptoms of urinary tract infections and treat promptly, if indicated [see *Adverse Reactions* (6)].

**5.6 Hypoglycemia with Concomitant Use with Insulin and Insulin Secretagogues**

Insulin and insulin secretagogues are known to cause hypoglycemia. INVOKANA can increase the risk of hypoglycemia when combined with insulin or an insulin secretagogue [see *Adverse Reactions* (6.1)]. Therefore, a lower dose of insulin or insulin secretagogue may be required to minimize the risk of hypoglycemia when used in combination with INVOKANA.

**5.7 Genital Mycotic Infections**

INVOKANA increases the risk of genital mycotic infections. Patients with a history of genital mycotic infections and uncircumcised males were more likely to develop genital mycotic infections [see *Adverse Reactions* (6.1)]. Monitor and treat appropriately.

**INVOKANA® (canagliflozin) tablets****5.8 Hypersensitivity Reactions**

Hypersensitivity reactions, including angioedema and anaphylaxis, have been reported with INVOKANA. These reactions generally occurred within hours to days after initiating INVOKANA. If hypersensitivity reactions occur, discontinue use of INVOKANA; treat and monitor until signs and symptoms resolve [see Contraindications (4) and Adverse Reactions (6.1, 6.2)].

**5.9 Bone Fracture**

An increased risk of bone fracture, occurring as early as 12 weeks after treatment initiation, was observed in patients using INVOKANA. Consider factors that contribute to fracture risk prior to initiating INVOKANA [see Adverse Reactions (6.1)].

**5.10 Increases in Low-Density Lipoprotein (LDL-C)**

Dose-related increases in LDL-C occur with INVOKANA [see Adverse Reactions (6.1)]. Monitor LDL-C and treat if appropriate after initiating INVOKANA.

**5.11 Macrovascular Outcomes**

There have been no clinical studies establishing conclusive evidence of macrovascular risk reduction with INVOKANA or any other antidiabetic drug.

**6 ADVERSE REACTIONS**

The following important adverse reactions are described below and elsewhere in the labeling:

- Hypotension [see Warnings and Precautions (5.1)]
- Ketoacidosis [see Warnings and Precautions (5.2)]
- Acute Kidney Injury and Impairment in Renal Function [see Warnings and Precautions (5.3)]
- Hyperkalemia [see Warnings and Precautions (5.4)]
- Urinary tract Infections [see Warnings and Precautions (5.5)]
- Hypoglycemia with Concomitant Use with Insulin and Insulin Secretagogues [see Warnings and Precautions (5.6)]
- Genital Mycotic Infections [see Warnings and Precautions (5.7)]
- Hypersensitivity Reactions [see Warnings and Precautions (5.8)]
- Bone Fracture [see Warnings and Precautions (5.9)]
- Increases in Low-Density Lipoprotein (LDL-C) [see Warnings and Precautions (5.10)]

**6.1 Clinical Studies Experience**

Because clinical trials are conducted under widely varying conditions, adverse reaction rates observed in the clinical trials of a drug cannot be directly compared to the rates in the clinical trials of another drug and may not reflect the rates observed in clinical practice.

**Pool of Placebo-Controlled Trials**

The data in Table 1 is derived from four 26-week placebo-controlled trials. In one trial INVOKANA was used as monotherapy and in three trials INVOKANA was used as add-on therapy [see Clinical Studies (14)]. These data reflect exposure of 1667 patients to INVOKANA and a mean duration of exposure to INVOKANA of 24 weeks. Patients received INVOKANA 100 mg (N=833), INVOKANA 300 mg (N=834) or placebo (N=646) once daily. The mean age of the population was 56 years and 2% were older than 75 years of age. Fifty percent (50%) of the population was male and 72% were Caucasian, 12% were Asian, and 5% were Black or African American. At baseline the population had diabetes for an average of 7.3 years, had a mean HbA1C of 8.0% and 20% had established microvascular complications of diabetes. Baseline renal function was normal or mildly impaired (mean eGFR 88 mL/min/1.73 m<sup>2</sup>).

Table 1 shows common adverse reactions associated with the use of INVOKANA. These adverse reactions were not present at baseline, occurred more commonly on INVOKANA than on placebo, and occurred in at least 2% of patients treated with either INVOKANA 100 mg or INVOKANA 300 mg.

**Table 1: Adverse Reactions From Pool of Four 26-Week Placebo-Controlled Studies Reported in ≥ 2% of INVOKANA-Treated Patients<sup>2</sup>**

Adverse Reaction	Placebo N=646	INVOKANA 100 mg N=833	INVOKANA 300 mg N=834
Urinary tract infections <sup>2</sup>	3.8%	6.9%	4.4%
Increased urination <sup>3</sup>	0.7%	5.1%	4.6%
Thirst <sup>4</sup>	0.1%	2.8%	2.4%
Constipation	0.8%	1.8%	2.4%
Nausea	1.6%	2.1%	2.3%
	N=312	N=425	N=430
Female genital mycotic infections <sup>1</sup>	2.8%	10.6%	11.6%
Vulvovaginal pruritus	0.0%	1.6%	3.2%
	N=334	N=408	N=404
Male genital mycotic infections <sup>1</sup>	0.7%	4.2%	3.8%

**INVOKANA® (canagliflozin) tablets**

\* The four placebo-controlled trials included one monotherapy trial and three add-on combination trials with metformin, metformin and sulfonylurea, or metformin and pioglitazone.

<sup>1</sup> Female genital mycotic infections include the following adverse reactions: Vulvovaginal candidiasis, Vulvovaginal mycotic infection, Vulvovaginitis, Vaginal infection, Vulvitis, and Genital infection fungal.

<sup>2</sup> Urinary tract Infections include the following adverse reactions: Urinary tract infection, Cystitis, Kidney infection, and Urosepsis.

<sup>3</sup> Increased urination includes the following adverse reactions: Polyuria, Pollakiuria, Urine output increased, Micturition urgency, and Nocturia.

<sup>4</sup> Male genital mycotic infections include the following adverse reactions: Balanitis or Balanoposthitis, Balanitis candida, and Genital infection fungal.

<sup>5</sup> Thirst includes the following adverse reactions: Thirst, Dry mouth, and Polydipsia.

Note: Percentages were weighted by studies. Study weights were proportional to the harmonic mean of the three treatment sample sizes.

Abdominal pain was also more commonly reported in patients taking INVOKANA 100 mg (1.8%), 300 mg (1.7%) than in patients taking placebo (0.8%).

**Pool of Placebo- and Active-Controlled Trials**

The occurrence of adverse reactions for canagliflozin was evaluated in a larger pool of patients participating in placebo- and active-controlled trials.

The data combined eight clinical trials [see Clinical Studies (14)] and reflect exposure of 6177 patients to INVOKANA. The mean duration of exposure to INVOKANA was 36 weeks with 1832 individuals exposed to INVOKANA for greater than 50 weeks. Patients received INVOKANA 100 mg (N=3092), INVOKANA 300 mg (N=3085) or comparator (N=3262) once daily. The mean age of the population was 60 years and 5% were older than 75 years of age. Fifty-eight percent (58%) of the population was male and 73% were Caucasian, 16% were Asian, and 4% were Black or African American. At baseline, the population had diabetes for an average of 11 years, had a mean HbA1C of 8.0% and 33% had established microvascular complications of diabetes. Baseline renal function was normal or mildly impaired (mean eGFR 81 mL/min/1.73 m<sup>2</sup>).

The types and frequency of common adverse reactions observed in the pool of eight clinical trials were consistent with those listed in Table 1. Percentages were weighted by studies. Study weights were proportional to the harmonic mean of the three treatment sample sizes. In this pool, INVOKANA was also associated with the adverse reactions of fatigue (1.8% with comparator, 2.2% with INVOKANA 100 mg, and 2.0% with INVOKANA 300 mg) and loss of strength or energy (i.e., asthenia) (0.6% with comparator, 0.7% with INVOKANA 100 mg, and 1.1% with INVOKANA 300 mg).

In the pool of eight clinical trials, the incidence rate of pancreatitis (acute or chronic) was 0.1, 0.2, and 0.1 receiving comparator, INVOKANA 100 mg, and INVOKANA 300 mg, respectively.

In the pool of eight clinical trials, hypersensitivity-related adverse reactions (including erythema, rash, pruritus, urticaria, and angioedema) occurred in 3.0%, 3.8%, and 4.2% of patients receiving comparator, INVOKANA 100 mg, and INVOKANA 300 mg, respectively. Five patients experienced serious adverse reactions of hypersensitivity with INVOKANA, which included 4 patients with urticaria and 1 patient with a diffuse rash and urticaria occurring within hours of exposure to INVOKANA. Among these patients, 2 patients discontinued INVOKANA. One patient with urticaria had recurrence when INVOKANA was re-initiated.

Photosensitivity-related adverse reactions (including photosensitivity reaction, polymorphic light eruption, and sunburn) occurred in 0.1%, 0.2%, and 0.2% of patients receiving comparator, INVOKANA 100 mg, and INVOKANA 300 mg, respectively.

Other adverse reactions occurring more frequently on INVOKANA than on comparator were:

**Volume Depletion-Related Adverse Reactions**

INVOKANA results in an osmotic diuresis, which may lead to reductions in intravascular volume. In clinical studies, treatment with INVOKANA was associated with a dose-dependent increase in the incidence of volume depletion-related adverse reactions (e.g., hypotension, postural dizziness, orthostatic hypotension, syncope, and dehydration). An increased incidence was observed in patients on the 300 mg dose. The three factors associated with the largest increase in volume depletion-related adverse reactions were the use of loop diuretics, moderate renal impairment (eGFR 30 to less than 60 mL/min/1.73 m<sup>2</sup>), and age 75 years and older (Table 2) [see Dosage and Administration (2.2), Warnings and Precautions (5.1), and Use in Specific Populations (8.5 and 8.6)].

**INVOKANA® (canagliflozin) tablets****Table 2: Proportion of Patients With at Least One Volume Depletion-Related Adverse Reaction (Pooled Results from 8 Clinical Trials)**

Baseline Characteristic	Comparator Group* %	INVOKANA 100 mg %	INVOKANA 300 mg %
Overall population	1.5%	2.3%	3.4%
75 years of age and older†	2.8%	4.9%	8.7%
eGFR less than 60 mL/min/1.73 m <sup>2</sup> ‡	2.5%	4.7%	8.1%
Use of loop diuretic†	4.7%	3.2%	8.8%

\* Includes placebo and active-comparator groups

† Patients could have more than 1 of the listed risk factors

**Falls**

In a pool of nine clinical trials with mean duration of exposure to INVOKANA of 85 weeks, the proportion of patients who experienced falls was 1.3%, 1.5%, and 2.1% with comparator, INVOKANA 100 mg, and INVOKANA 300 mg, respectively. The higher risk of falls for patients treated with INVOKANA was observed within the first few weeks of treatment.

**Impairment in Renal Function**

INVOKANA is associated with a dose-dependent increase in serum creatinine and a concomitant fall in estimated GFR (Table 3). Patients with moderate renal impairment at baseline had larger mean changes.

**Table 3: Changes in Serum Creatinine and eGFR Associated with INVOKANA in the Pool of Four Placebo-Controlled Trials and Moderate Renal Impairment Trial**

			Placebo	INVOKANA	INVOKANA
			N=646	100 mg N=833	300 mg N=834
Pool of Four Placebo-Controlled Trials	Baseline	Creatinine (mg/dL)	0.84	0.82	0.82
		eGFR (mL/min/1.73 m <sup>2</sup> )	87.0	88.3	88.8
	Week 8 Change	Creatinine (mg/dL)	0.01	0.03	0.05
		eGFR (mL/min/1.73 m <sup>2</sup> )	-1.6	-3.8	-5.0
	End of Treatment Change*	Creatinine (mg/dL)	0.01	0.02	0.03
		eGFR (mL/min/1.73 m <sup>2</sup> )	-1.6	-2.3	-3.4
Moderate Renal Impairment Trial	Baseline	Creatinine (mg/dL)	1.61	1.62	1.63
		eGFR (mL/min/1.73 m <sup>2</sup> )	40.1	39.7	36.5
	Week 3 Change	Creatinine (mg/dL)	0.03	0.18	0.28
		eGFR (mL/min/1.73 m <sup>2</sup> )	-0.7	-4.6	-8.2
	End of Treatment Change*	Creatinine (mg/dL)	0.07	0.18	0.18
		eGFR (mL/min/1.73 m <sup>2</sup> )	-1.5	-3.8	-4.0

\* Week 28 in MITT LOCF population

In the pool of four placebo-controlled trials where patients had normal or mildly impaired baseline renal function, the proportion of patients who experienced at least one event of significant renal function decline, defined as an eGFR below 80 mL/min/1.73 m<sup>2</sup> and 30% lower than baseline, was 2.1% with placebo, 2.0% with INVOKANA 100 mg, and 4.1% with INVOKANA 300 mg. At the end of treatment, 0.5% with placebo, 0.7% with INVOKANA 100 mg, and 1.4% with INVOKANA 300 mg had a significant renal function decline.

In a trial carried out in patients with moderate renal impairment with a baseline eGFR of 30 to less than 50 mL/min/1.73 m<sup>2</sup> (mean baseline eGFR 39 mL/min/1.73 m<sup>2</sup>) [see Clinical Studies (14.3)], the proportion of patients who experienced at least one event of significant renal function decline, defined as an eGFR 30% lower than baseline, was 6.9% with placebo, 18% with INVOKANA 100 mg, and 22.5% with INVOKANA 300 mg. At the end of treatment, 4.6% with placebo, 3.4% with INVOKANA 100 mg, and 2.2% with INVOKANA 300 mg had a significant renal function decline.

In a pooled population of patients with moderate renal impairment (N=1085) with baseline eGFR of 30 to less than 60 mL/min/1.73 m<sup>2</sup> (mean baseline eGFR 48 mL/min/1.73 m<sup>2</sup>), the overall incidence of these events was lower than in the dedicated trial but a dose-dependent increase in incident episodes of significant renal function decline compared to placebo was still observed.

Use of INVOKANA has been associated with an increased incidence of renal-related adverse reactions (e.g., increased blood creatinine, decreased glomerular filtration rate, renal impairment, and acute renal failure), particularly in patients with moderate renal impairment.

**INVOKANA® (canagliflozin) tablets**

In the pooled analysis of patients with moderate renal impairment, the incidence of renal-related adverse reactions was 3.7% with placebo, 8.8% with INVOKANA 100 mg, and 9.3% with INVOKANA 300 mg. Discontinuations due to renal-related adverse events occurred in 1.0% with placebo, 1.2% with INVOKANA 100 mg, and 1.8% with INVOKANA 300 mg [see Warnings and Precautions (5.3)].

**Genital Mycotic Infections**

In the pool of four placebo-controlled clinical trials, female genital mycotic infections (e.g., vulvovaginal mycotic infection, vulvovaginal candidiasis, and vulvovaginitis) occurred in 2.8%, 10.6%, and 11.6% of females treated with placebo, INVOKANA 100 mg, and INVOKANA 300 mg, respectively. Patients with a history of genital mycotic infections were more likely to develop genital mycotic infections on INVOKANA. Female patients who developed genital mycotic infections on INVOKANA were more likely to experience recurrence and require treatment with oral or topical antifungal agents and anti-microbial agents. In females, discontinuation due to genital mycotic infections occurred in 0% and 0.7% of patients treated with placebo and INVOKANA, respectively [see Warnings and Precautions (5.7)].

In the pool of four placebo-controlled clinical trials, male genital mycotic infections (e.g., candidal balanitis, balanoposthitis) occurred in 0.7%, 4.2%, and 3.8% of males treated with placebo, INVOKANA 100 mg, and INVOKANA 300 mg, respectively. Male genital mycotic infections occurred more commonly in uncircumcised males and in males with a prior history of balanitis or balanoposthitis. Male patients who developed genital mycotic infections on INVOKANA were more likely to experience recurrent infections (22% on INVOKANA versus none on placebo), and require treatment with oral or topical antifungal agents and anti-microbial agents than patients on comparators. In males, discontinuations due to genital mycotic infections occurred in 0% and 0.5% of patients treated with placebo and INVOKANA, respectively. In the pooled analysis of 8 controlled trials, phimosis was reported in 0.3% of uncircumcised male patients treated with INVOKANA and 0.2% required circumcision to treat the phimosis [see Warnings and Precautions (5.7)].

**Hypoglycemia**

In all clinical trials, hypoglycemia was defined as any event regardless of symptoms, where biochemical hypoglycemia was documented (any glucose value below or equal to 70 mg/dL). Severe hypoglycemia was defined as an event consistent with hypoglycemia where the patient required the assistance of another person to recover, lost consciousness, or experienced a seizure (regardless of whether biochemical documentation of a low glucose value was obtained). In individual clinical trials [see Clinical Studies (14)], episodes of hypoglycemia occurred at a higher rate when INVOKANA was co-administered with insulin or sulfonylureas (Table 4) [see Warnings and Precautions (5.6)].

**Table 4: Incidence of Hypoglycemia\* in Controlled Clinical Studies**

Monotherapy (26 weeks)	Placebo (N=182)	INVOKANA 100 mg (N=185)	INVOKANA 300 mg (N=197)
Overall [N (%)]	5 (2.8)	7 (3.8)	8 (3.0)
In Combination with Metformin (26 weeks)	Placebo + Metformin (N=183)	INVOKANA 100 mg + Metformin (N=366)	INVOKANA 300 mg + Metformin (N=367)
Overall [N (%)]	3 (1.8)	16 (4.3)	17 (4.6)
Severe [N (%)]†	0 (0)	1 (0.3)	1 (0.3)
In Combination with Metformin (52 weeks)	Glimepiride + Metformin (N=482)	INVOKANA 100 mg + Metformin (N=483)	INVOKANA 300 mg + Metformin (N=485)
Overall [N (%)]	165 (34.2)	27 (5.6)	24 (4.9)
Severe [N (%)]†	15 (3.1)	2 (0.4)	3 (0.6)
In Combination with Sulfonylurea (18 weeks)	Placebo + Sulfonylurea (N=69)	INVOKANA 100 mg + Sulfonylurea (N=74)	INVOKANA 300 mg + Sulfonylurea (N=72)
Overall [N (%)]	4 (5.8)	3 (4.1)	8 (12.5)
In Combination with Metformin + Sulfonylurea (26 weeks)	Placebo + Metformin + Sulfonylurea (N=156)	INVOKANA 100 mg + Metformin + Sulfonylurea (N=157)	INVOKANA 300 mg + Metformin + Sulfonylurea (N=156)
Overall [N (%)]	24 (16.4)	43 (27.4)	47 (30.1)
Severe [N (%)]†	1 (0.6)	1 (0.6)	0
In Combination with Metformin + Sulfonylurea (52 weeks)	Sitagliptin + Metformin + Sulfonylurea (N=378)		INVOKANA 300 mg + Metformin + Sulfonylurea (N=377)
Overall [N (%)]	154 (40.7)		183 (43.2)
Severe [N (%)]†	13 (3.4)		15 (4.0)

**INVOKANA® (canagliflozin) tablets****Table 4: Incidence of Hypoglycemia\* in Controlled Clinical Studies (continued)**

In Combination with Metformin + Pioglitazone (26 weeks)	Placebo + Metformin + Pioglitazone (N=115)	INVOKANA 100 mg + Metformin + Pioglitazone (N=113)	INVOKANA 300 mg + Metformin + Pioglitazone (N=114)
Overall [N (%)]	3 (2.6)	3 (2.7)	6 (5.3)
In Combination with Insulin (18 weeks)	Placebo (N=585)	INVOKANA 100 mg (N=566)	INVOKANA 300 mg (N=587)
Overall [N (%)]	208 (36.8)	279 (49.3)	285 (48.6)
Severe [N (%)]†	14 (2.5)	10 (1.9)	16 (2.7)

\* Number of patients experiencing at least one event of hypoglycemia based on either biochemically documented episodes or severe hypoglycemic events in the intent-to-treat population

† Severe episodes of hypoglycemia were defined as those where the patient required the assistance of another person to recover, lost consciousness, or experienced a seizure (regardless of whether biochemical documentation of a low glucose value was obtained)

**Bone Fracture**

The occurrence of bone fractures was evaluated in a pool of nine clinical trials with a mean duration of exposure to INVOKANA of 85 weeks. The incidence rates of adjudicated bone fractures were 1.1, 1.4, and 1.5 per 100 patient-years of exposure in the comparator, INVOKANA 100 mg, and INVOKANA 300 mg groups, respectively. Fractures were observed as early as 12 weeks after treatment initiation and were more likely to be low trauma (e.g., fall from no more than standing height), and affect the upper extremities [see Warnings and Precautions (5.9)].

**Laboratory and Imaging Tests****Increases in Serum Potassium**

In a pooled population of patients (N=723) with moderate renal impairment (eGFR 45 to less than 60 mL/min/1.73 m<sup>2</sup>), increases in serum potassium to greater than 5.4 mEq/L and 15% above baseline occurred in 5.3%, 6.0%, and 8.8% of patients treated with placebo, INVOKANA 100 mg, and INVOKANA 300 mg, respectively. Severe elevations (greater than or equal to 6.5 mEq/L) occurred in 0.4% of patients treated with placebo, no patients treated with INVOKANA 100 mg, and 1.3% of patients treated with INVOKANA 300 mg.

In these patients, increases in potassium were more commonly seen in those with elevated potassium at baseline. Among patients with moderate renal impairment, approximately 84% were taking medications that interfere with potassium excretion, such as potassium-sparing diuretics, angiotensin-converting-enzyme inhibitors, and angiotensin-receptor blockers [see Warnings and Precautions (5.3 and 5.4) and Use in Specific Populations (8.6)].

**Increases in Serum Magnesium**

Dose-related increases in serum magnesium were observed early after initiation of INVOKANA (within 6 weeks) and remained elevated throughout treatment. In the pool of four placebo-controlled trials, the mean percent change in serum magnesium levels was 8.1% and 9.3% with INVOKANA 100 mg and INVOKANA 300 mg, respectively, compared to -0.6% with placebo. In a trial of patients with moderate renal impairment [see Clinical Studies (14.3)], serum magnesium levels increased by 0.2%, 8.2%, and 14.8% with placebo, INVOKANA 100 mg, and INVOKANA 300 mg, respectively.

**Increases in Serum Phosphate**

Dose-related increases in serum phosphate levels were observed with INVOKANA. In the pool of four placebo-controlled trials, the mean percent change in serum phosphate levels were 3.8% and 5.1% with INVOKANA 100 mg and INVOKANA 300 mg, respectively, compared to 1.5% with placebo. In a trial of patients with moderate renal impairment [see Clinical Studies (14.3)], the mean serum phosphate levels increased by 1.2%, 6.0%, and 9.3% with placebo, INVOKANA 100 mg, and INVOKANA 300 mg, respectively.

**Increases in Low-Density Lipoprotein Cholesterol (LDL-C) and non-High-Density Lipoprotein Cholesterol (non-HDL-C)**

In the pool of four placebo-controlled trials, dose-related increases in LDL-C with INVOKANA were observed. Mean changes (percent changes) from baseline in LDL-C relative to placebo were 4.4 mg/dL (4.5%) and 9.2 mg/dL (8.0%) with INVOKANA 100 mg and INVOKANA 300 mg, respectively. The mean baseline LDL-C levels were 104 to 110 mg/dL across treatment groups [see Warnings and Precautions (5.10)].

Dose-related increases in non-HDL-C with INVOKANA were observed. Mean changes (percent changes) from baseline in non-HDL-C relative to placebo were 2.1 mg/dL (1.5%) and 5.1 mg/dL (3.6%) with INVOKANA 100 mg and 300 mg, respectively. The mean baseline non-HDL-C levels were 140 to 147 mg/dL across treatment groups.

**INVOKANA® (canagliflozin) tablets****Increases in Hemoglobin**

In the pool of four placebo-controlled trials, mean changes (percent changes) from baseline in hemoglobin were -0.18 g/dL (-1.1%) with placebo, 0.47 g/dL (3.5%) with INVOKANA 100 mg, and 0.51 g/dL (3.8%) with INVOKANA 300 mg. The mean baseline hemoglobin value was approximately 14.1 g/dL across treatment groups. At the end of treatment, 0.8%, 4.0%, and 2.7% of patients treated with placebo, INVOKANA 100 mg, and INVOKANA 300 mg, respectively, had hemoglobin above the upper limit of normal.

**Decreases in Bone Mineral Density**

Bone mineral density (BMD) was measured by dual-energy X-ray absorptiometry in a clinical trial of 714 older adults (mean age 64 years) [see Clinical Studies (14.3)]. At 2 years, patients randomized to INVOKANA 100 mg and INVOKANA 300 mg had placebo-corrected declines in BMD at the total hip of 0.9% and 1.2%, respectively, and at the lumbar spine of 0.3% and 0.7%, respectively. Additionally, placebo-adjusted BMD declines were 0.1% at the femoral neck for both INVOKANA doses and 0.4% at the distal forearm for patients randomized to INVOKANA 300 mg. The placebo-adjusted change at the distal forearm for patients randomized to INVOKANA 100 mg was 0%.

**6.2 Postmarketing Experience**

Additional adverse reactions have been identified during postapproval use of INVOKANA. Because these reactions are reported voluntarily from a population of uncertain size, it is generally not possible to reliably estimate their frequency or establish a causal relationship to drug exposure.

**Ketoacidosis [see Warnings and Precautions (5.2)]**

Acute Kidney Injury and Impairment in Renal Function [see Warnings and Precautions (5.3)]

Anaphylaxis, Angioedema [see Warnings and Precautions (5.8)]

Urosepsis and Pyelonephritis [see Warnings and Precautions (5.5)]

**7 DRUG INTERACTIONS****7.1 UGT Enzyme Inducers**

Rifampin: Co-administration of canagliflozin with rifampin, a nonselective inducer of several UGT enzymes, including UGT1A9, UGT2B4, decreased canagliflozin area under the curve (AUC) by 51%. This decrease in exposure to canagliflozin may decrease efficacy. If an inducer of these UGTs (e.g., rifampin, phenytoin, phenobarbital, ritonavir) must be co-administered with INVOKANA (canagliflozin), consider increasing the dose to 300 mg once daily if patients are currently tolerating INVOKANA 100 mg once daily, have an eGFR greater than 60 mL/min/1.73 m<sup>2</sup>, and require additional glycemic control. Consider other antihyperglycemic therapy in patients with an eGFR of 45 to less than 60 mL/min/1.73 m<sup>2</sup> receiving concurrent therapy with a UGT inducer and require additional glycemic control [see Dosage and Administration (2.3) and Clinical Pharmacology (12.3)].

**7.2 Digoxin**

There was an increase in the AUC and mean peak drug concentration (C<sub>max</sub>) of digoxin (20% and 36%, respectively) when co-administered with INVOKANA 300 mg [see Clinical Pharmacology (12.3)]. Patients taking INVOKANA with concomitant digoxin should be monitored appropriately.

**7.3 Positive Urine Glucose Test**

Monitoring glycemic control with urine glucose tests is not recommended in patients taking SGLT2 inhibitors as SGLT2 inhibitors increase urinary glucose excretion and will lead to positive urine glucose tests. Use alternative methods to monitor glycemic control.

**7.4 Interference with 1,5-anhydroglucitol (1,5-AG) Assay**

Monitoring glycemic control with 1,5-AG assay is not recommended as measurements of 1,5-AG are unreliable in assessing glycemic control in patients taking SGLT2 inhibitors. Use alternative methods to monitor glycemic control.

**8 USE IN SPECIFIC POPULATIONS****8.1 Pregnancy****Risk Summary**

Based on animal data showing adverse renal effects, INVOKANA is not recommended during the second and third trimesters of pregnancy.

Limited data with INVOKANA in pregnant women are not sufficient to determine a drug-associated risk for major birth defects or miscarriage. There are risks to the mother and fetus associated with poorly controlled diabetes in pregnancy [see Clinical Considerations].

In animal studies, adverse renal pelvic and tubule dilatations that were not reversible were observed in rats when canagliflozin was administered during a period of renal development corresponding to the late second and third trimesters of human pregnancy, at an exposure 0.5-times the 300 mg clinical dose, based on AUC.

**INVOKANA® (canagliflozin) tablets**

The estimated background risk of major birth defects is 6-10% in women with pre-gestational diabetes with a HbA1c >7 and has been reported to be as high as 20-25% in women with a HbA1c >10. The estimated background risk of miscarriage for the indicated population is unknown. In the U.S. general population, the estimated background risk of major birth defects and miscarriage in clinically recognized pregnancies is 2-4% and 15-20%, respectively.

**Clinical Considerations****Disease-associated maternal and/or embryo/fetal risk**

Poorly controlled diabetes in pregnancy increases the maternal risk for diabetic ketoacidosis, pre-eclampsia, spontaneous abortions, preterm delivery, stillbirth and delivery complications. Poorly controlled diabetes increases the fetal risk for major birth defects, stillbirth, and macrosomia related morbidity.

**Animal Data**

Canagliflozin dosed directly to juvenile rats from postnatal day (PND) 21 until PND 90 at doses of 4, 20, 65, or 100 mg/kg increased kidney weights and dose dependently increased the incidence and severity of renal pelvic and tubular dilatation at all doses tested. Exposure at the lowest dose was greater than or equal to 0.5-times the 300 mg clinical dose, based on AUC. These outcomes occurred with drug exposure during periods of renal development in rats that correspond to the late second and third trimester of human renal development. The renal pelvic dilatations observed in juvenile animals did not fully reverse within a 1 month recovery period.

In embryo-fetal development studies in rats and rabbits, canagliflozin was administered for intervals coinciding with the first trimester period of organogenesis in humans. No developmental toxicities independent of maternal toxicity were observed when canagliflozin was administered at doses up to 100 mg/kg in pregnant rats and 160 mg/kg in pregnant rabbits during embryonic organogenesis or during a study in which maternal rats were dosed from gestation day (GD) 6 through PND 21, yielding exposures up to approximately 19-times the 300 mg clinical dose, based on AUC.

**8.2 Lactation****Risk Summary**

There is no information regarding the presence of INVOKANA in human milk, the effects on the breastfed infant, or the effects on milk production. Canagliflozin is present in the milk of lactating rats [see Data]. Since human kidney maturation occurs *in utero* and during the first 2 years of life when lactational exposure may occur, there may be risk to the developing human kidney.

Because of the potential for serious adverse reactions in a breastfed infant, advise women that use of INVOKANA is not recommended while breastfeeding.

**Data****Animal Data**

Rediolabeled canagliflozin administered to lactating rats on day 13 post-partum was present at a milk/plasma ratio of 1.40, indicating that canagliflozin and its metabolites are transferred into milk at a concentration comparable to that in plasma. Juvenile rats directly exposed to canagliflozin showed a risk to the developing kidney (renal pelvic and tubular dilatations) during maturation.

**8.4 Pediatric Use**

Safety and effectiveness of INVOKANA in pediatric patients under 18 years of age have not been established.

**8.5 Geriatric Use**

Two thousand thirty-four (2034) patients 65 years and older, and 345 patients 75 years and older were exposed to INVOKANA in nine clinical studies of INVOKANA [see Clinical Studies (14.3)].

Patients 65 years and older had a higher incidence of adverse reactions related to reduced intravascular volume with INVOKANA (such as hypotension, postural dizziness, orthostatic hypotension, syncope, and dehydration), particularly with the 300 mg daily dose, compared to younger patients; a more prominent increase in the incidence was seen in patients who were 75 years and older [see Dosage and Administration (2.1) and Adverse Reactions (6.1)]. Smaller reductions in HbA1C with INVOKANA relative to placebo were seen in older (65 years and older; -0.81% with INVOKANA 100 mg and -0.74% with INVOKANA 300 mg relative to placebo) compared to younger patients (-0.72% with INVOKANA 100 mg and -0.87% with INVOKANA 300 mg relative to placebo).

**8.6 Renal Impairment**

The efficacy and safety of INVOKANA were evaluated in a study that included patients with moderate renal impairment (eGFR 30 to less than 50 mL/min/1.73 m<sup>2</sup>) [see Clinical Studies (14.3)]. These patients had less

**INVOKANA® (canagliflozin) tablets**

overall glycemic efficacy and had a higher occurrence of adverse reactions related to reduced intravascular volume, renal-related adverse reactions, and decreases in eGFR compared to patients with mild renal impairment or normal renal function (eGFR greater than or equal to 60 mL/min/1.73 m<sup>2</sup>). Dose-related, transient mean increases in serum potassium were observed early after initiation of INVOKANA (i.e., within 3 weeks) in this trial. Increases in serum potassium of greater than 5.4 mEq/L and 15% above baseline occurred in 16.1%, 12.4%, and 27.0% of patients treated with placebo, INVOKANA 100 mg, and INVOKANA 300 mg, respectively. Severe elevations (greater than or equal to 6.5 mEq/L) occurred in 1.1%, 2.2%, and 2.2% of patients treated with placebo, INVOKANA 100 mg, and INVOKANA 300 mg, respectively [see Dosage and Administration (2.2), Warnings and Precautions (5.1, 5.3, and 5.4), and Adverse Reactions (6.1)].

The efficacy and safety of INVOKANA have not been established in patients with severe renal impairment (eGFR less than 30 mL/min/1.73 m<sup>2</sup>), with ESRD, or receiving dialysis. INVOKANA is not expected to be effective in these patient populations [see Contraindications (4) and Clinical Pharmacology (12.3)].

**8.7 Hepatic Impairment**

No dosage adjustment is necessary in patients with mild or moderate hepatic impairment. The use of INVOKANA has not been studied in patients with severe hepatic impairment and is therefore not recommended [see Clinical Pharmacology (12.3)].


**10 OVERDOSAGE**

There were no reports of overdose during the clinical development program of INVOKANA (canagliflozin).

In the event of an overdose, contact the Poison Control Center. It is also reasonable to employ the usual supportive measures, e.g., remove unabsorbed material from the gastrointestinal tract, employ clinical monitoring, and institute supportive treatment as dictated by the patient's clinical status. Canagliflozin was negligibly removed during a 4-hour hemodialysis session. Canagliflozin is not expected to be dialyzable by peritoneal dialysis.

**11 DESCRIPTION**

INVOKANA (canagliflozin) contains canagliflozin, an inhibitor of sodium-glucose co-transporter 2 (SGLT2), the transporter responsible for reabsorbing the majority of glucose filtered by the kidney. Canagliflozin, the active ingredient of INVOKANA, is chemically known as (1S)-1,5-anhydro-1-[3-[[5-(4-fluorophenyl)-2-thienyl]methyl]-4-methylphenyl]-D-glucitol hemihydrate and its molecular formula and weight are C<sub>24</sub>H<sub>29</sub>FO<sub>5</sub>S•1/2 H<sub>2</sub>O and 453.53, respectively. The structural formula for canagliflozin is:



Canagliflozin is practically insoluble in aqueous media from pH 1.1 to 12.9.

INVOKANA is supplied as film-coated tablets for oral administration, containing 102 and 306 mg of canagliflozin in each tablet strength, corresponding to 100 mg and 300 mg of canagliflozin (anhydrous), respectively.

Inactive ingredients of the core tablet are croscarmellose sodium, hydroxypropyl cellulose, lactose anhydrous, magnesium stearate, and microcrystalline cellulose. The magnesium stearate is vegetable-sourced. The tablets are finished with a commercially available film-coating consisting of the following excipients: polyvinyl alcohol (partially hydrolyzed), titanium dioxide, macrogol/PEG, talc, and iron oxide yellow, E172 (100 mg tablet only).

**12 CLINICAL PHARMACOLOGY****12.1 Mechanism of Action**

Sodium-glucose co-transporter 2 (SGLT2), expressed in the proximal renal tubules, is responsible for the majority of the reabsorption of filtered glucose from the tubular lumen. Canagliflozin is an inhibitor of SGLT2. By inhibiting SGLT2, canagliflozin reduces reabsorption of filtered glucose and lowers the renal threshold for glucose (RT<sub>G</sub>), and thereby increases urinary glucose excretion (UGE).

**INVOKANA® (canagliflozin) tablets****12.2 Pharmacodynamics**

Following single and multiple oral doses of canagliflozin in patients with type 2 diabetes, dose-dependent decreases in the renal threshold for glucose (RT<sub>G</sub>) and increases in urinary glucose excretion were observed. From a starting RT<sub>G</sub> value of approximately 240 mg/dL, canagliflozin at 100 mg and 300 mg once daily suppressed RT<sub>G</sub> throughout the 24-hour period. Maximal suppression of mean RT<sub>G</sub> over the 24-hour period was seen with the 300 mg daily dose to approximately 70 to 80 mg/dL in patients with type 2 diabetes in Phase 1 studies. The reductions in RT<sub>G</sub> led to increases in mean UGE of approximately 100 g/day in subjects with type 2 diabetes treated with either 100 mg or 300 mg of canagliflozin. In patients with type 2 diabetes given 100 mg to 300 mg once daily over a 16-day dosing period, reductions in RT<sub>G</sub> and increases in urinary glucose excretion were observed over the dosing period. In this study, plasma glucose declined in a dose-dependent fashion within the first day of dosing. In single-dose studies in healthy and type 2 diabetic subjects, treatment with canagliflozin 300 mg before a mixed-meal delayed intestinal glucose absorption and reduced postprandial glucose.

**Cardiac Electrophysiology**

In a randomized, double-blind, placebo-controlled, active-comparator, 4-way crossover study, 60 healthy subjects were administered a single oral dose of canagliflozin 300 mg, canagliflozin 1,200 mg (4 times the maximum recommended dose), moxifloxacin, and placebo. No meaningful changes in QTc interval were observed with either the recommended dose of 300 mg or the 1,200 mg dose.

**12.3 Pharmacokinetics**

The pharmacokinetics of canagliflozin is similar in healthy subjects and patients with type 2 diabetes. Following single-dose oral administration of 100 mg and 300 mg of INVOKANA, peak plasma concentrations (median T<sub>max</sub>) of canagliflozin occurs within 1 to 2 hours post-dose. Plasma C<sub>max</sub> and AUC of canagliflozin increased in a dose-proportional manner from 50 mg to 300 mg. The apparent terminal half-life (t<sub>1/2</sub>) was 10.8 hours and 13.1 hours for the 100 mg and 300 mg doses, respectively. Steady-state was reached after 4 to 5 days of once-daily dosing with canagliflozin 100 mg to 300 mg. Canagliflozin does not exhibit time-dependent pharmacokinetics and accumulated in plasma up to 36% following multiple doses of 100 mg and 300 mg.

**Absorption**

The mean absolute oral bioavailability of canagliflozin is approximately 65%. Co-administration of a high-fat meal with canagliflozin had no effect on the pharmacokinetics of canagliflozin; therefore, INVOKANA may be taken with or without food. However, based on the potential to reduce postprandial plasma glucose excursions due to delayed intestinal glucose absorption, it is recommended that INVOKANA be taken before the first meal of the day [see *Dosage and Administration* (2.1)].

**Distribution**

The mean steady-state volume of distribution of canagliflozin following a single intravenous infusion in healthy subjects was 83.5 L, suggesting extensive tissue distribution. Canagliflozin is extensively bound to proteins in plasma (99%), mainly to albumin. Protein binding is independent of canagliflozin plasma concentrations. Plasma protein binding is not meaningfully altered in patients with renal or hepatic impairment.

**Metabolism**

O-glucuronidation is the major metabolic elimination pathway for canagliflozin, which is mainly glucuronidated by UGT1A9 and UGT2B4 to two inactive O-glucuronide metabolites.

CYP3A4-mediated (oxidative) metabolism of canagliflozin is minimal (approximately 7%) in humans.

**Excretion**

Following administration of a single oral [<sup>14</sup>C] canagliflozin dose to healthy subjects, 41.5%, 7.0%, and 3.2% of the administered radioactive dose was recovered in feces as canagliflozin, a hydroxylated metabolite, and an O-glucuronide metabolite, respectively. Enterohepatic circulation of canagliflozin was negligible.

Approximately 33% of the administered radioactive dose was excreted in urine, mainly as O-glucuronide metabolites (30.5%). Less than 1% of the dose was excreted as unchanged canagliflozin in urine. Renal clearance of canagliflozin 100 mg and 300 mg doses ranged from 1.30 to 1.55 mL/min.

Mean systemic clearance of canagliflozin was approximately 182 mL/min in healthy subjects following intravenous administration.

**Specific Populations****Renal Impairment**

A single-dose, open-label study evaluated the pharmacokinetics of canagliflozin 200 mg in subjects with varying degrees of renal impairment (classified using the MDRD-eGFR formula) compared to healthy subjects.

**INVOKANA® (canagliflozin) tablets**

Renal impairment did not affect the C<sub>max</sub> of canagliflozin. Compared to healthy subjects (N=3; eGFR greater than or equal to 90 mL/min/1.73 m<sup>2</sup>), plasma AUC of canagliflozin was increased by approximately 15%, 29%, and 53% in subjects with mild (N=10), moderate (N=9), and severe (N=10) renal impairment, respectively, (eGFR 60 to less than 90, 30 to less than 60 and 15 to less than 30 mL/min/1.73 m<sup>2</sup>, respectively), but was similar for ESRD (N=8) subjects and healthy subjects.

Increases in canagliflozin AUC of this magnitude are not considered clinically relevant. The pharmacodynamic response to canagliflozin declines with increasing severity of renal impairment [see *Contraindications* (4) and *Warnings and Precautions* (5.3)].

Canagliflozin was negligibly removed by hemodialysis.

**Hepatic Impairment**

Relative to subjects with normal hepatic function, the geometric mean ratios for C<sub>max</sub> and AUC<sub>0-∞</sub> of canagliflozin were 107% and 110%, respectively, in subjects with Child-Pugh class A (mild hepatic impairment) and 96% and 111%, respectively, in subjects with Child-Pugh class B (moderate hepatic impairment) following administration of a single 300 mg dose of canagliflozin.

These differences are not considered to be clinically meaningful. There is no clinical experience in patients with Child-Pugh class C (severe) hepatic impairment [see *Use in Specific Populations* (8.7)].

**Pharmacokinetic Effects of Age, Body Mass Index (BMI)/Weight, Gender and Race**

Based on the population PK analysis with data collected from 1526 subjects, age, body mass index (BMI)/weight, gender, and race do not have a clinically meaningful effect on the pharmacokinetics of canagliflozin [see *Use in Specific Populations* (8.5)].

**Pediatric**

Studies characterizing the pharmacokinetics of canagliflozin in pediatric patients have not been conducted.

**Drug Interaction Studies****In Vitro Assessment of Drug Interactions**

Canagliflozin did not induce CYP450 enzyme expression (3A4, 2C9, 2C19, 2B6, and 1A2) in cultured human hepatocytes. Canagliflozin did not inhibit the CYP450 isoenzymes (1A2, 2A6, 2C19, 2D6, or 2E1) and weakly inhibited CYP2B6, CYP2C8, CYP2C9, and CYP3A4 based on *in vitro* studies with human hepatic microsomes. Canagliflozin is a weak inhibitor of P-gp.

Canagliflozin is also a substrate of drug transporters P-glycoprotein (P-gp) and MRP2.

**In Vivo Assessment of Drug Interactions**

**Table 5: Effect of Co-Administered Drugs on Systemic Exposures of Canagliflozin**

Co-Administered Drug	Dose of Co-Administered Drug*	Dose of Canagliflozin*	Geometric Mean Ratio (Ratio With/Without Co-Administered Drug) No Effect=1.0	
			AUC <sup>†</sup> (90% CI)	C <sub>max</sub> (90% CI)
<b>See Drug Interactions (7.1) for the clinical relevance of the following:</b>				
Rifampin	600 mg QD for 8 days	300 mg	0.49 (0.44; 0.54)	0.72 (0.61; 0.84)
<b>No dose adjustments of INVOKANA required for the following:</b>				
Cyclosporine	400 mg	300 mg QD for 8 days	1.23 (1.19; 1.27)	1.01 (0.91; 1.11)
Ethinyl estradiol and levonorgestrel	0.03 mg ethinyl estradiol and 0.15 mg levonorgestrel	200 mg QD for 8 days	0.91 (0.88; 0.94)	0.92 (0.84; 0.99)
Hydrochlorothiazide	25 mg QD for 35 days	300 mg QD for 7 days	1.12 (1.08; 1.17)	1.15 (1.06; 1.25)
Metformin	2,000 mg	300 mg QD for 8 days	1.10 (1.05; 1.16)	1.05 (0.98; 1.16)
Probeneclid	500 mg BID for 3 days	300 mg QD for 17 days	1.21 (1.16; 1.25)	1.13 (1.00; 1.28)

\* Single dose unless otherwise noted

<sup>†</sup> AUC<sub>inf</sub> for drugs given as a single dose and AUC<sub>24h</sub> for drugs given as multiple doses

QD = once daily; BID = twice daily

**INVOKANA® (canagliflozin) tablets****Table 6: Effect of Canagliflozin on Systemic Exposure of Co-Administered Drugs**

Co-Administered Drug	Dose of Co-Administered Drug <sup>†</sup>	Dose of Canagliflozin <sup>†</sup>	Geometric Mean Ratio (Ratio With/Without Co-Administered Drug) No Effect = 1.0	AUC <sup>‡</sup>		C <sub>max</sub>	
				(90% CI)	(90% CI)	(90% CI)	(90% CI)
See <i>Drug Interactions (7.2)</i> for the clinical relevance of the following:							
Digoxin	0.5 mg QD first day followed by 0.25 mg QD for 6 days	300 mg QD for 7 days	digoxin	1.20 (1.12; 1.28)	1.36 (1.21; 1.53)		
No dose adjustments of co-administered drug required for the following:							
Acetaminophen	1,000 mg	300 mg BID for 25 days	acetaminophen	1.05 <sup>†</sup> (0.98; 1.14)	1.00 (0.92; 1.09)		
Ethinyl estradiol and levonorgestrel	0.03 mg ethinyl estradiol and 0.15 mg levonorgestrel	200 mg QD for 6 days	ethinyl estradiol	1.07 (0.99; 1.15)	1.22 (1.10; 1.36)		
			levonorgestrel	1.06 (1.00; 1.13)	1.22 (1.11; 1.35)		
Glyburide	1.25 mg	200 mg QD for 6 days	glyburide	1.02 (0.98; 1.07)	0.93 (0.85; 1.01)		
			3-cis-hydroxy-glyburide	1.01 (0.96; 1.07)	0.99 (0.91; 1.08)		
			4-trans-hydroxy-glyburide	1.03 (0.97; 1.09)	0.96 (0.88; 1.04)		
Hydrochlorothiazide	25 mg QD for 35 days	300 mg QD for 7 days	hydrochlorothiazide	0.99 (0.95; 1.04)	0.94 (0.87; 1.01)		
Metformin	2,000 mg	300 mg QD for 6 days	metformin	1.20 (1.08; 1.34)	1.06 (0.93; 1.20)		
Simvastatin	40 mg	300 mg QD for 7 days	simvastatin	1.12 (0.94; 1.33)	1.09 (0.91; 1.31)		
			simvastatin acid	1.18 (1.03; 1.35)	1.26 (1.10; 1.45)		
Warfarin	30 mg	300 mg QD for 12 days	(R)-warfarin	1.01 (0.96; 1.06)	1.03 (0.94; 1.13)		
			(S)-warfarin	1.06 (1.00; 1.12)	1.01 (0.93; 1.13)		
			INR	1.00 (0.96; 1.03)	1.05 (0.99; 1.12)		

<sup>†</sup> Single dose unless otherwise noted

<sup>‡</sup> AUC<sub>inf</sub> for drugs given as a single dose and AUC<sub>0-12h</sub> for drugs given as multiple doses

<sup>†</sup> AUC<sub>0-12h</sub>

QD = once daily; BID = twice daily; INR = International Normalized Ratio

**13 NONCLINICAL TOXICOLOGY****13.1 Carcinogenesis, Mutagenesis, Impairment of Fertility****Carcinogenesis**

Carcinogenicity was evaluated in 2-year studies conducted in CD1 mice and Sprague-Dawley rats. Canagliflozin did not increase the incidence of tumors in mice dosed at 10, 30, or 100 mg/kg (less than or equal to 14 times exposure from a 300 mg clinical dose).

Testicular Leydig cell tumors, considered secondary to increased luteinizing hormone (LH), increased significantly in male rats at all doses tested (10, 30, and 100 mg/kg). In a 12-week clinical study, LH did not increase in males treated with canagliflozin.

Renal tubular adenoma and carcinoma increased significantly in male and female rats dosed at 100 mg/kg, or approximately 12-times exposure from a 300 mg clinical dose. Also, adrenal pheochromocytoma increased significantly in males and numerically in females dosed at 100 mg/kg. Carbohydrate malabsorption associated with high doses of canagliflozin was considered a necessary proximal event in the emergence of renal and adrenal tumors in rats. Clinical studies have not demonstrated carbohydrate malabsorption in humans at canagliflozin doses of up to 2-times the recommended clinical dose of 300 mg.

**Mutagenesis**

Canagliflozin was not mutagenic with or without metabolic activation in the Ames assay. Canagliflozin was mutagenic in the *in vitro* mouse lymphoma assay with but not without metabolic activation. Canagliflozin was not mutagenic or clastogenic in an *in vivo* oral micronucleus assay in rats and an *in vivo* oral Comet assay in rats.

**Impairment of Fertility**

Canagliflozin had no effects on the ability of rats to mate and sire or maintain a litter up to the high dose of 100 mg/kg (approximately 14 times and 18 times the 300 mg clinical dose in males and females, respectively), although there were minor alterations in a number of reproductive parameters (decreased

**INVOKANA® (canagliflozin) tablets**

sperm velocity, increased number of abnormal sperm, slightly fewer corpora lutea, fewer implantation sites, and smaller litter sizes) at the highest dosage administered.

**14 CLINICAL STUDIES**

INVOKANA (canagliflozin) has been studied as monotherapy, in combination with metformin, sulfonylurea, metformin and sulfonylurea, metformin and a thiazolidinedione (i.e., pioglitazone), and in combination with insulin (with or without other antihyperglycemic agents). The efficacy of INVOKANA was compared to a dipeptidyl peptidase-4 (DPP-4) inhibitor (sitagliptin) and a sulfonylurea (glimepiride). INVOKANA was also evaluated in adults 55 to 80 years of age and patients with moderate renal impairment.

In patients with type 2 diabetes, treatment with INVOKANA produced clinically and statistically significant improvements in HbA1C compared to placebo. Reductions in HbA1C were observed across subgroups including age, gender, race, and baseline body mass index (BMI).

**14.1 Monotherapy**

A total of 584 patients with type 2 diabetes inadequately controlled on diet and exercise participated in a 26-week, double-blind, placebo-controlled study to evaluate the efficacy and safety of INVOKANA. The mean age was 55 years, 44% of patients were men, and the mean baseline eGFR was 87 mL/min/1.73 m<sup>2</sup>. Patients taking other antihyperglycemic agents (N=281) discontinued the agent and underwent an 8-week washout followed by a 2-week, single-blind, placebo run-in period. Patients not taking oral antihyperglycemic agents (N=303) entered the 2-week, single-blind, placebo run-in period directly. After the placebo run-in period, patients were randomized to INVOKANA 100 mg, INVOKANA 300 mg, or placebo, administered once daily for 28 weeks.

At the end of treatment, INVOKANA 100 mg and 300 mg once daily resulted in a statistically significant improvement in HbA1C (p<0.001 for both doses) compared to placebo. INVOKANA 100 mg and 300 mg once daily also resulted in a greater proportion of patients achieving an HbA1C less than 7%, in significant reduction in fasting plasma glucose (FPG), in improved postprandial glucose (PPG), and in percent body weight reduction compared to placebo (see Table 7). Statistically significant (p<0.001 for both doses) mean changes from baseline in systolic blood pressure relative to placebo were -3.7 mmHg and -5.4 mmHg with INVOKANA 100 mg and 300 mg, respectively.

**Table 7: Results from 28-Week Placebo-Controlled Clinical Study with INVOKANA as Monotherapy<sup>‡</sup>**

Efficacy Parameter	Placebo (N=192)	INVOKANA 100 mg (N=195)	INVOKANA 300 mg (N=197)
<b>HbA1C (%)</b>			
Baseline (mean)	7.97	8.06	8.01
Change from baseline (adjusted mean)	0.14	-0.77	-1.03
Difference from placebo (adjusted mean) (95% CI) <sup>†</sup>		-0.91 <sup>‡</sup> (-1.09; -0.73)	-1.16 <sup>‡</sup> (-1.34; -0.99)
Percent of Patients Achieving HbA1C < 7%	21	45 <sup>‡</sup>	62 <sup>‡</sup>
<b>Fasting Plasma Glucose (mg/dL)</b>			
Baseline (mean)	188	172	173
Change from baseline (adjusted mean)	8	-27	-35
Difference from placebo (adjusted mean) (95% CI) <sup>†</sup>		-36 <sup>‡</sup> (-42; -29)	-43 <sup>‡</sup> (-50; -37)
<b>2-hour Postprandial Glucose (mg/dL)</b>			
Baseline (mean)	229	250	254
Change from baseline (adjusted mean)	5	-43	-59
Difference from placebo (adjusted mean) (95% CI) <sup>†</sup>		-48 <sup>‡</sup> (-59.1; -37.0)	-64 <sup>‡</sup> (-75.0; -52.9)
<b>Body Weight</b>			
Baseline (mean) in kg	87.5	85.9	86.9
% change from baseline (adjusted mean)	-0.6	-2.8	-3.9
Difference from placebo (adjusted mean) (95% CI) <sup>†</sup>		-2.2 <sup>‡</sup> (-2.9; -1.6)	-3.3 <sup>‡</sup> (-4.0; -2.6)

<sup>‡</sup> Intent-to-treat population using last observation in study prior to glycemic rescue therapy

<sup>†</sup> Least squares mean adjusted for baseline value and stratification factors

<sup>‡</sup> p<0.001

**INVOKANA® (canagliflozin) tablets****14.2 Combination Therapy****Add-on Combination Therapy with Metformin**

A total of 1284 patients with type 2 diabetes inadequately controlled on metformin monotherapy (greater than or equal to 2,000 mg/day, or at least 1,500 mg/day if higher dose not tolerated) participated in a 26-week, double-blind, placebo- and active-controlled study to evaluate the efficacy and safety of INVOKANA in combination with metformin. The mean age was 55 years, 47% of patients were men, and the mean baseline eGFR was 89 mL/min/1.73 m<sup>2</sup>. Patients already on the required metformin dose (N=1009) were randomized after completing a 2-week, single-blind, placebo run-in period. Patients taking less than the required metformin dose or patients on metformin in combination with another antihyperglycemic agent (N=275) were switched to metformin monotherapy (at doses described above) for at least 8 weeks before entering the 2-week, single-blind, placebo run-in. After the placebo run-in period, patients were randomized to INVOKANA 100 mg, INVOKANA 300 mg, sitagliptin 100 mg, or placebo, administered once daily as add-on therapy to metformin.

At the end of treatment, INVOKANA 100 mg and 300 mg once daily resulted in a statistically significant improvement in HbA1C (p<0.001 for both doses) compared to placebo when added to metformin. INVOKANA 100 mg and 300 mg once daily also resulted in a greater proportion of patients achieving an HbA1C less than 7%, in significant reduction in fasting plasma glucose (FPG), in improved postprandial glucose (PPG), and in percent body weight reduction compared to placebo when added to metformin (see Table 8). Statistically significant (p<0.001 for both doses) mean changes from baseline in systolic blood pressure relative to placebo were -5.4 mmHg and -6.6 mmHg with INVOKANA 100 mg and 300 mg, respectively.

**Table 8: Results from 26-Week Placebo-Controlled Clinical Study of INVOKANA In Combination with Metformin\***

Efficacy Parameter	Placebo + Metformin (N=163)	INVOKANA 100 mg + Metformin (N=368)	INVOKANA 300 mg + Metformin (N=367)
<b>HbA1C (%)</b>			
Baseline (mean)	7.96	7.94	7.95
Change from baseline (adjusted mean)	-0.17	-0.79	-0.94
Difference from placebo (adjusted mean) (95% CI) <sup>†</sup>		-0.62 <sup>‡</sup> (-0.78; -0.48)	-0.77 <sup>‡</sup> (-0.91; -0.64)
Percent of patients achieving HbA1C < 7%	30	46 <sup>‡</sup>	58 <sup>‡</sup>
<b>Fasting Plasma Glucose (mg/dL)</b>			
Baseline (mean)	164	169	173
Change from baseline (adjusted mean)	2	-27	-38
Difference from placebo (adjusted mean) (95% CI) <sup>†</sup>		-30 <sup>‡</sup> (-36; -24)	-40 <sup>‡</sup> (-46; -34)
<b>2-hour Postprandial Glucose (mg/dL)</b>			
Baseline (mean)	249	258	262
Change from baseline (adjusted mean)	-10	-48	-57
Difference from placebo (adjusted mean) (95% CI) <sup>†</sup>		-38 <sup>‡</sup> (-49; -27)	-47 <sup>‡</sup> (-58; -36)
<b>Body Weight</b>			
Baseline (mean) in kg	86.7	88.7	85.4
% change from baseline (adjusted mean)	-1.2	-3.7	-4.2
Difference from placebo (adjusted mean) (95% CI) <sup>†</sup>		-2.5 <sup>‡</sup> (-3.1; -1.9)	-2.9 <sup>‡</sup> (-3.5; -2.3)

\* Intent-to-treat population using last observation in study prior to glycemic rescue therapy

<sup>†</sup> Least squares mean adjusted for baseline value and stratification factors

<sup>‡</sup> p<0.001

**Initial Combination Therapy with Metformin**

A total of 1186 patients with type 2 diabetes inadequately controlled with diet and exercise participated in a 26-week double-blind, active-controlled, parallel-group, 5-arm, multicenter study to evaluate the efficacy and safety of initial therapy with INVOKANA in combination with metformin XR. The median

**INVOKANA® (canagliflozin) tablets**

age was 56 years, 48% of patients were men, and the mean baseline eGFR was 87.8 mL/min/1.73 m<sup>2</sup>. The median duration of diabetes was 1.8 years, and 72% of patients were treatment naïve. After completing a 2-week single-blind placebo run-in period, patients were randomly assigned for a double-blind treatment period of 26 weeks to 1 of 5 treatment groups (Table 9). The metformin XR dose was initiated at 500 mg/day for the first week of treatment and then increased to 1000 mg/day. Metformin XR or matching placebo was up-titrated every 2-3 weeks during the next 6 weeks of treatment to a maximum daily dose of 1500 to 2000 mg/day, as tolerated; about 90% of patients reached 2000 mg/day.

At the end of treatment, INVOKANA 100 mg and INVOKANA 300 mg in combination with metformin XR resulted in a statistically significant greater improvement in HbA1C compared to their respective INVOKANA doses (100 mg and 300 mg) alone or metformin XR alone.

**Table 9: Results from 26-Week Active-Controlled Clinical Study of INVOKANA Alone or INVOKANA as Initial Combination Therapy with Metformin\***

Efficacy Parameter	Metformin XR (N=237)	INVOKANA 100 mg (N=237)	INVOKANA 300 mg (N=238)	INVOKANA 100 mg + Metformin XR (N=237)	INVOKANA 300 mg + Metformin XR (N=237)
<b>HbA1C (%)</b>					
Baseline (mean)	8.81	8.78	8.77	8.83	8.80
Change from baseline (adjusted mean) <sup>†</sup>	-1.30	-1.37	-1.42	-1.77	-1.78
Difference from canagliflozin 100 mg (adjusted mean) (95% CI) <sup>‡</sup>				-0.40 <sup>‡</sup> (-0.59; -0.21)	
Difference from canagliflozin 300 mg (adjusted mean) (95% CI) <sup>‡</sup>					-0.35 <sup>‡</sup> (-0.56; -0.17)
Difference from metformin XR (adjusted mean) (95% CI) <sup>‡</sup>		-0.06 <sup>††</sup> (-0.26; 0.13)	-0.11 <sup>††</sup> (-0.31; 0.08)	-0.48 <sup>†</sup> (-0.66; -0.27)	-0.48 <sup>†</sup> (-0.67; -0.28)
Percent of patients achieving HbA1C < 7%	38	34	39	47 <sup>‡‡</sup>	51 <sup>‡‡</sup>

\* Intent-to-treat population

<sup>†</sup> Least squares mean adjusted for covariates including baseline value and stratification factor

<sup>‡</sup> Adjusted p=0.001 for superiority

<sup>‡‡</sup> Adjusted p=0.001 for non-inferiority

<sup>‡‡‡</sup> Adjusted p<0.05

<sup>†††</sup> There were 121 patients without week 26 efficacy data. Analysis addressing missing data gave consistent results with the results provided in this table.

**INVOKANA Compared to Glimepiride, Both as Add-on Combination With Metformin**

A total of 1450 patients with type 2 diabetes inadequately controlled on metformin monotherapy (greater than or equal to 2,000 mg/day, or at least 1,500 mg/day if higher dose not tolerated) participated in a 52-week, double-blind, active-controlled study to evaluate the efficacy and safety of INVOKANA in combination with metformin.

The mean age was 58 years, 52% of patients were men, and the mean baseline eGFR was 80 mL/min/1.73 m<sup>2</sup>. Patients tolerating maximally required metformin dose (N=928) were randomized after completing a 2-week, single-blind, placebo run-in period. Other patients (N=522) were switched to metformin monotherapy (at doses described above) for at least 10 weeks, then completed a 2-week single-blind run-in period. After the 2-week run-in period, patients were randomized to INVOKANA 100 mg, INVOKANA 300 mg, or glimepiride (titration allowed throughout the 52-week study to 6 or 8 mg), administered once daily as add-on therapy to metformin.

As shown in Table 10 and Figure 1, at the end of treatment, INVOKANA 100 mg provided similar reductions in HbA1C from baseline compared to glimepiride when added to metformin therapy. INVOKANA 300 mg provided a greater reduction from baseline in HbA1C compared to glimepiride, and the relative treatment difference was -0.12% (95% CI: -0.22; -0.02). As shown in Table 10, treatment with INVOKANA 100 mg and 300 mg daily provided greater improvements in percent body weight change, relative to glimepiride.


**INVOKANA® (canagliflozin) tablets**

**Table 10: Results from 52-Week Clinical Study Comparing INVOKANA to Glimepiride in Combination with Metformin\***

Efficacy Parameter	INVOKANA 100 mg + Metformin (N=483)	INVOKANA 300 mg + Metformin (N=485)	Glimepiride (titrated) + Metformin (N=482)
<b>HbA1C (%)</b>			
Baseline (mean)	7.78	7.79	7.83
Change from baseline (adjusted mean)	-0.82	-0.93	-0.81
Difference from glimepiride (adjusted mean) (95% CI) <sup>†</sup>	-0.01 <sup>‡</sup> (-0.11; 0.09)	-0.12 <sup>‡</sup> (-0.22; -0.02)	
Percent of patients achieving HbA1C < 7%	54	60	56
<b>Fasting Plasma Glucose (mg/dL)</b>			
Baseline (mean)	185	184	188
Change from baseline (adjusted mean)	-24	-28	-18
Difference from glimepiride (adjusted mean) (95% CI) <sup>‡</sup>	-6 (-10; -2)	-9 (-13; -5)	
<b>Body Weight</b>			
Baseline (mean) in kg	86.8	86.6	88.8
% change from baseline (adjusted mean)	-4.2	-4.7	1.0
Difference from glimepiride (adjusted mean) (95% CI) <sup>‡</sup>	-5.2 <sup>‡</sup> (-5.7; -4.7)	-5.7 <sup>‡</sup> (-6.2; -5.1)	

\* Intent-to-treat population using last observation in study prior to glycemic rescue therapy  
<sup>†</sup> Least squares mean adjusted for baseline value and stratification factors  
<sup>‡</sup> INVOKANA + metformin is considered non-inferior to glimepiride + metformin because the upper limit of this confidence interval is less than the pre-specified non-inferiority margin of < 0.3%.  
<sup>§</sup> p<0.001

**Figure 1: Mean HbA1C Change at Each Time Point (Completers) and at Week 52 Using Last Observation Carried Forward (mITT Population)**



**Add-on Combination Therapy With Sulfonylurea**

A total of 127 patients with type 2 diabetes inadequately controlled on sulfonylurea monotherapy participated in an 18-week, double-blind, placebo-controlled sub-study to evaluate the efficacy and safety of INVOKANA in combination with sulfonylurea. The mean age was 65 years, 57% of patients were men, and the mean baseline eGFR was 69 mL/min/1.73 m<sup>2</sup>. Patients treated with sulfonylurea monotherapy on a stable protocol-specified dose (greater than or equal to 50% maximal dose) for at least 10 weeks completed a 2-week, single-blind, placebo run-in period. During the run-in period, patients with inadequate glycemic control were randomized to INVOKANA 100 mg, INVOKANA 300 mg, or placebo, administered once daily as add-on to sulfonylurea.

As shown in Table 11, at the end of treatment, INVOKANA 100 mg and 300 mg daily provided statistically significant (p<0.001 for both doses) improvements in HbA1C relative to placebo when added to sulfonylurea. INVOKANA 300 mg once daily compared to placebo resulted in a greater proportion of patients achieving an HbA1C less than 7%, (33% vs 5%), greater reductions in fasting plasma glucose (-36 mg/dL vs +12 mg/dL), and greater percent body weight reduction (-2.0% vs -0.2%).

**INVOKANA® (canagliflozin) tablets**

**Table 11: Results from 18-Week Placebo-Controlled Clinical Study of INVOKANA in Combination with Sulfonylurea\***

Efficacy Parameter	Placebo + Sulfonylurea (N=45)	INVOKANA 100 mg + Sulfonylurea (N=42)	INVOKANA 300 mg + Sulfonylurea (N=40)
<b>HbA1C (%)</b>			
Baseline (mean)	8.49	8.29	8.28
Change from baseline (adjusted mean)	0.04	-0.70	-0.79
Difference from placebo (adjusted mean) (95% CI) <sup>†</sup>		-0.74 <sup>‡</sup> (-1.15; -0.33)	-0.83 <sup>‡</sup> (-1.24; -0.41)

\* Intent-to-treat population using last observation in study prior to glycemic rescue therapy  
<sup>†</sup> Least squares mean adjusted for baseline value  
<sup>‡</sup> p<0.001

**Add-on Combination Therapy With Metformin and Sulfonylurea**

A total of 489 patients with type 2 diabetes inadequately controlled on the combination of metformin (greater than or equal to 2,000 mg/day or at least 1,500 mg/day if higher dose not tolerated) and sulfonylurea (maximal or near-maximal effective dose) participated in a 26-week, double-blind, placebo-controlled study to evaluate the efficacy and safety of INVOKANA in combination with metformin and sulfonylurea. The mean age was 57 years, 51% of patients were men, and the mean baseline eGFR was 89 mL/min/1.73 m<sup>2</sup>. Patients already on the protocol-specified doses of metformin and sulfonylurea (N=372) entered a 2-week, single-blind, placebo run-in period. Other patients (N=97) were required to be on a stable protocol-specified dose of metformin and sulfonylurea for at least 8 weeks before entering the 2-week run-in period. Following the run-in period, patients were randomized to INVOKANA 100 mg, INVOKANA 300 mg, or placebo, administered once daily as add-on to metformin and sulfonylurea.

At the end of treatment, INVOKANA 100 mg and 300 mg once daily resulted in a statistically significant improvement in HbA1C (p<0.001 for both doses) compared to placebo when added to metformin and sulfonylurea. INVOKANA 100 mg and 300 mg once daily also resulted in a greater proportion of patients achieving an HbA1C less than 7%, in a significant reduction in fasting plasma glucose (FPG), and in percent body weight reduction compared to placebo when added to metformin and sulfonylurea (see Table 12).

**Table 12: Results from 26-Week Placebo-Controlled Clinical Study of INVOKANA in Combination with Metformin and Sulfonylurea\***

Efficacy Parameter	Placebo + Metformin and Sulfonylurea (N=156)	INVOKANA 100 mg + Metformin and Sulfonylurea (N=157)	INVOKANA 300 mg + Metformin and Sulfonylurea (N=156)
<b>HbA1C (%)</b>			
Baseline (mean)	8.12	8.13	8.13
Change from baseline (adjusted mean)	-0.13	-0.85	-1.06
Difference from placebo (adjusted mean) (95% CI) <sup>†</sup>		-0.71 <sup>‡</sup> (-0.90; -0.52)	-0.92 <sup>‡</sup> (-1.11; -0.73)
Percent of patients achieving A1C < 7%	18	43 <sup>‡</sup>	57 <sup>‡</sup>
<b>Fasting Plasma Glucose (mg/dL)</b>			
Baseline (mean)	170	173	168
Change from baseline (adjusted mean)	4	-18	-31
Difference from placebo (adjusted mean) (95% CI) <sup>‡</sup>		-22 <sup>‡</sup> (-31; -13)	-35 <sup>‡</sup> (-44; -25)
<b>Body Weight</b>			
Baseline (mean) in kg	90.8	93.5	93.5
% change from baseline (adjusted mean)	-0.7	-2.1	-2.6
Difference from placebo (adjusted mean) (95% CI) <sup>‡</sup>		-1.4 <sup>‡</sup> (-2.1; -0.7)	-2.0 <sup>‡</sup> (-2.7; -1.3)

\* Intent-to-treat population using last observation in study prior to glycemic rescue therapy  
<sup>†</sup> Least squares mean adjusted for baseline value and stratification factors  
<sup>‡</sup> p<0.001

**INVOKANA® (canagliflozin) tablets**

**INVOKANA Compared to Sitagliptin, Both as Add-on Combination Therapy With Metformin and Sulfonylurea**

A total of 755 patients with type 2 diabetes inadequately controlled on the combination of metformin (greater than or equal to 2,000 mg/day or at least 1,500 mg/day if higher dose not tolerated) and sulfonylurea (near-maximal or maximal effective dose) participated in a 52-week, double-blind, active-controlled study to compare the efficacy and safety of INVOKANA 300 mg versus sitagliptin 100 mg in combination with metformin and sulfonylurea. The mean age was 57 years, 56% of patients were men, and the mean baseline eGFR was 88 mL/min/1.73 m<sup>2</sup>. Patients already on protocol-specified doses of metformin and sulfonylurea (N=716) entered a 2-week single-blind, placebo run-in period. Other patients (N=39) were required to be on a stable protocol-specified dose of metformin and sulfonylurea for at least 8 weeks before entering the 2-week run-in period. Following the run-in period, patients were randomized to INVOKANA 300 mg or sitagliptin 100 mg as add-on to metformin and sulfonylurea.

As shown in Table 13 and Figure 2, at the end of treatment, INVOKANA 300 mg provided greater HbA1C reduction compared to sitagliptin 100 mg when added to metformin and sulfonylurea (p<0.05). INVOKANA 300 mg resulted in a mean percent change in body weight from baseline of -2.5% compared to +0.3% with sitagliptin 100 mg. A mean change in systolic blood pressure from baseline of -5.06 mmHg was observed with INVOKANA 300 mg compared to +0.85 mmHg with sitagliptin 100 mg.

**Table 13: Results from 52-Week Clinical Study Comparing INVOKANA to Sitagliptin in Combination with Metformin and Sulfonylurea\***

Efficacy Parameter	INVOKANA 300 mg + Metformin and Sulfonylurea (N=377)	Sitagliptin 100 mg + Metformin and Sulfonylurea (N=378)
<b>HbA1C (%)</b>		
Baseline (mean)	8.12	8.13
Change from baseline (adjusted mean)	-1.03	-0.68
Difference from sitagliptin (adjusted mean) (95% CI) <sup>†</sup>	-0.37 <sup>‡</sup>	
	(-0.50; -0.25)	
Percent of patients achieving HbA1C < 7%	48	35
<b>Fasting Plasma Glucose (mg/dL)</b>		
Baseline (mean)	170	164
Change from baseline (adjusted mean)	-30	-6
Difference from sitagliptin (adjusted mean) (95% CI) <sup>†</sup>	-24	
	(-30; -18)	
<b>Body Weight</b>		
Baseline (mean) in kg	87.6	89.6
% change from baseline (adjusted mean)	-2.5	0.3
Difference from sitagliptin (adjusted mean) (95% CI) <sup>†</sup>	-2.8 <sup>‡</sup>	
	(-3.3; -2.2)	


\* Intent-to-treat population using last observation in study prior to glycemic rescue therapy

<sup>†</sup> Least squares mean adjusted for baseline value and stratification factors

<sup>‡</sup> INVOKANA + metformin + sulfonylurea is considered non-inferior to sitagliptin + metformin + sulfonylurea because the upper limit of this confidence interval is less than the pre-specified non-inferiority margin of < 0.3%.

<sup>§</sup> p<0.001

**Figure 2: Mean HbA1C Change at Each Time Point (Completers) and at Week 52 Using Last Observation Carried Forward (mITT Population)**



**INVOKANA® (canagliflozin) tablets**

**Add-on Combination Therapy With Metformin and Pioglitazone**

A total of 342 patients with type 2 diabetes inadequately controlled on the combination of metformin (greater than or equal to 2,000 mg/day or at least 1,500 mg/day if higher dose not tolerated) and pioglitazone (30 or 45 mg/day) participated in a 26-week, double-blind, placebo-controlled study to evaluate the efficacy and safety of INVOKANA in combination with metformin and pioglitazone. The mean age was 57 years, 63% of patients were men, and the mean baseline eGFR was 86 mL/min/1.73 m<sup>2</sup>. Patients already on protocol-specified doses of metformin and pioglitazone (N=163) entered a 2-week, single-blind, placebo run-in period. Other patients (N=181) were required to be on stable protocol-specified doses of metformin and pioglitazone for at least 8 weeks before entering the 2-week run-in period. Following the run-in period, patients were randomized to INVOKANA 100 mg, INVOKANA 300 mg, or placebo, administered once daily as add-on to metformin and pioglitazone.

At the end of treatment, INVOKANA 100 mg and 300 mg once daily resulted in a statistically significant improvement in HbA1C (p<0.001 for both doses) compared to placebo when added to metformin and pioglitazone. INVOKANA 100 mg and 300 mg once daily also resulted in a greater proportion of patients achieving an HbA1C less than 7%, in significant reduction in fasting plasma glucose (FPG) and in percent body weight reduction compared to placebo when added to metformin and pioglitazone (see Table 14). Statistically significant (p<0.05 for both doses) mean changes from baseline in systolic blood pressure relative to placebo were -4.1 mmHg and -3.5 mmHg with INVOKANA 100 mg and 300 mg, respectively.

**Table 14: Results from 26-Week Placebo-Controlled Clinical Study of INVOKANA in Combination with Metformin and Pioglitazone\***

Efficacy Parameter	Placebo + Metformin and Pioglitazone (N=115)	INVOKANA 100 mg + Metformin and Pioglitazone (N=113)	INVOKANA 300 mg + Metformin and Pioglitazone (N=114)
<b>HbA1C (%)</b>			
Baseline (mean)	8.00	7.99	7.84
Change from baseline (adjusted mean)	-0.26	-0.89	-1.03
Difference from placebo (adjusted mean) (95% CI) <sup>†</sup>		-0.62 <sup>‡</sup>	-0.76 <sup>‡</sup>
		(-0.81; -0.44)	(-0.95; -0.58)
Percent of patients achieving HbA1C < 7%	33	47 <sup>‡</sup>	64 <sup>‡</sup>
<b>Fasting Plasma Glucose (mg/dL)</b>			
Baseline (mean)	164	169	164
Change from baseline (adjusted mean)	3	-27	-33
Difference from placebo (adjusted mean) (95% CI) <sup>†</sup>		-29 <sup>‡</sup>	-36 <sup>‡</sup>
		(-37; -22)	(-43; -28)
<b>Body Weight</b>			
Baseline (mean) in kg	94.0	94.2	94.4
% change from baseline (adjusted mean)	-0.1	-2.8	-3.8
Difference from placebo (adjusted mean) (95% CI) <sup>†</sup>		-2.7 <sup>‡</sup>	-3.7 <sup>‡</sup>
		(-3.6; -1.8)	(-4.6; -2.8)

\* Intent-to-treat population using last observation in study prior to glycemic rescue therapy

<sup>†</sup> Least squares mean adjusted for baseline value and stratification factors

<sup>‡</sup> p<0.001

**Add-On Combination Therapy With Insulin (With or Without Other Antihyperglycemic Agents)**

A total of 1719 patients with type 2 diabetes inadequately controlled on insulin greater than or equal to 30 units/day or insulin in combination with other antihyperglycemic agents participated in an 18-week, double-blind, placebo-controlled substudy of a cardiovascular study to evaluate the efficacy and safety of INVOKANA in combination with insulin. The mean age was 63 years, 66% of patients were men, and the mean baseline eGFR was 75 mL/min/1.73 m<sup>2</sup>. Patients on basal, bolus, or basal/bolus insulin for at least 10 weeks entered a 2-week, single-blind, placebo run-in period. Approximately 70% of patients were on a background basal/bolus insulin regimen. After the run-in period, patients were randomized to INVOKANA 100 mg, INVOKANA 300 mg, or placebo, administered once daily as add-on to insulin. The mean daily insulin dose at baseline was 83 units, which was similar across treatment groups.

**INVOKANA® (canagliflozin) tablets**

At the end of treatment, INVOKANA 100 mg and 300 mg once daily resulted in a statistically significant improvement in HbA1C ( $p < 0.001$  for both doses) compared to placebo when added to insulin. INVOKANA 100 mg and 300 mg once daily also resulted in a greater proportion of patients achieving an HbA1C less than 7%, in significant reductions in fasting plasma glucose (FPG), and in percent body weight reductions compared to placebo (see Table 15). Statistically significant ( $p < 0.001$  for both doses) mean changes from baseline in systolic blood pressure relative to placebo were -2.8 mmHg and -4.4 mmHg with INVOKANA 100 mg and 300 mg, respectively.

**Table 15: Results from 18-Week Placebo-Controlled Clinical Study of INVOKANA in Combination with Insulin  $\geq$  30 Units/Day (With or Without Other Oral Antihyperglycemic Agents)\***

Efficacy Parameter	Placebo + Insulin (N=565)	INVOKANA 100 mg + Insulin (N=566)	INVOKANA 300 mg + Insulin (N=587)
<b>HbA1C (%)</b>			
Baseline (mean)	8.20	8.33	8.27
Change from baseline (adjusted mean)	0.01	-0.63	-0.72
Difference from placebo (adjusted mean) (95% CI)†		-0.65‡ (-0.73; -0.56)	-0.73‡ (-0.82; -0.65)
Percent of patients achieving HbA1C < 7%	8	20‡	25‡
<b>Fasting Plasma Glucose (mg/dL)</b>			
Baseline	169	170	168
Change from baseline (adjusted mean)	4	-19	-25
Difference from placebo (adjusted mean) (97.5% CI)‡		-23‡ (-29; -16)	-29‡ (-35; -23)
<b>Body Weight</b>			
Baseline (mean) in kg	97.7	96.9	96.7
% change from baseline (adjusted mean)	0.1	-1.8	-2.3
Difference from placebo (adjusted mean) (97.5% CI)‡		-1.9‡ (-2.2; -1.6)	-2.4‡ (-2.7; -2.1)

\* Intent-to-treat population using last observation in study prior to glycemic rescue therapy

† Least squares mean adjusted for baseline value and stratification factors

‡  $p < 0.001$

### 14.3 Studies in Special Populations

#### Adults 55 to 80 Years of Age

A total of 714 older patients with type 2 diabetes inadequately controlled on current diabetes therapy (either diet and exercise alone or in combination with oral or parenteral agents) participated in a 26-week, double-blind, placebo-controlled study to evaluate the efficacy and safety of INVOKANA in combination with current diabetes treatment. The mean age was 64 years, 55% of patients were men, and the mean baseline eGFR was 77 mL/min/1.73 m<sup>2</sup>. Patients were randomized to the addition of INVOKANA 100 mg, INVOKANA 300 mg, or placebo, administered once daily. At the end of treatment, INVOKANA provided statistically significant improvements from baseline relative to placebo in HbA1C ( $p < 0.001$  for both doses) of -0.57% (95% CI: -0.71; -0.44) for INVOKANA 100 mg and -0.70% (95% CI: -0.84; -0.57) for INVOKANA 300 mg. Statistically significant ( $p < 0.001$  for both doses) reductions from baseline in fasting plasma glucose (FPG) and body weight were also observed in this study relative to placebo [see Use in Specific Populations (8.5)].

#### Moderate Renal Impairment

A total of 269 patients with type 2 diabetes and a baseline eGFR of 30 mL/min/1.73 m<sup>2</sup> to less than 50 mL/min/1.73 m<sup>2</sup> inadequately controlled on current diabetes therapy participated in a 26-week, double-blind, placebo-controlled clinical study to evaluate the efficacy and safety of INVOKANA in combination with current diabetes treatment (diet or antihyperglycemic agent therapy, with 95% of patients on insulin and/or sulfonylureas). The mean age was 60 years, 61% of patients were men, and the mean baseline eGFR was 39 mL/min/1.73 m<sup>2</sup>. Patients were randomized to the addition of INVOKANA 100 mg, INVOKANA 300 mg, or placebo, administered once daily. At the end of treatment, INVOKANA 100 mg and INVOKANA 300 mg daily provided greater reductions in HbA1C relative to placebo (-0.30% [95% CI: -0.53; -0.07] and -0.40%, [95% CI: -0.64; -0.17], respectively) [see Warnings and Precautions (5.3), Adverse Reactions (6.1), and Use in Specific Populations (8.6)].

**INVOKANA® (canagliflozin) tablets**

### 16 HOW SUPPLIED/STORAGE AND HANDLING

INVOKANA (canagliflozin) tablets are available in the strengths and packages listed below:

100 mg tablets are yellow, capsule-shaped, film-coated tablets with "CFZ" on one side and "100" on the other side.

NDC 50458-140-30	Bottle of 30
NDC 50458-140-90	Bottle of 90
NDC 50458-140-50	Bottle of 500
NDC 50458-140-10	Blister package containing 100 tablets (10 blister cards containing 10 tablets each)

300 mg tablets are white, capsule-shaped, film-coated tablets with "CFZ" on one side and "300" on the other side.

NDC 50458-141-30	Bottle of 30
NDC 50458-141-90	Bottle of 90
NDC 50458-141-50	Bottle of 500
NDC 50458-141-10	Blister package containing 100 tablets (10 blister cards containing 10 tablets each)

#### Storage and Handling

Store at 25°C (77°F); excursions permitted to 15 to 30°C (59 to 86°F).

### 17 PATIENT COUNSELING INFORMATION

See FDA-approved patient labeling (Medication Guide).

#### Instructions

Instruct patients to read the Medication Guide before starting INVOKANA (canagliflozin) therapy and to reread it each time the prescription is renewed.

Inform patients of the potential risks and benefits of INVOKANA and of alternative modes of therapy. Also inform patients about the importance of adherence to dietary instructions, regular physical activity, periodic blood glucose monitoring and HbA1C testing, recognition and management of hypoglycemia and hyperglycemia, and assessment for diabetes complications. Advise patients to seek medical advice promptly during periods of stress such as fever, trauma, infection, or surgery, as medication requirements may change.

Instruct patients to take INVOKANA only as prescribed. If a dose is missed, advise patients to take it as soon as it is remembered unless it is almost time for the next dose, in which case patients should skip the missed dose and take the medicine at the next regularly scheduled time. Advise patients not to take two doses of INVOKANA at the same time.

Inform patients that the most common adverse reactions associated with INVOKANA are genital mycotic infection, urinary tract infection, and increased urination.

Inform female patients of child bearing age that the use of INVOKANA during pregnancy has not been studied in humans, and that INVOKANA should only be used during pregnancy only if the potential benefit justifies the potential risk to the fetus. Instruct patients to report pregnancies to their physicians as soon as possible.

Inform nursing mothers to discontinue INVOKANA or nursing, taking into account the importance of drug to the mother.

#### Laboratory Tests

Due to its mechanism of action, patients taking INVOKANA will test positive for glucose in their urine.

#### Hypotension

Inform patients that symptomatic hypotension may occur with INVOKANA and advise them to contact their doctor if they experience such symptoms [see Warnings and Precautions (5.1)]. Inform patients that dehydration may increase the risk for hypotension, and to have adequate fluid intake.

#### Ketoacidosis

Inform patients that ketoacidosis is a serious life-threatening condition. Cases of ketoacidosis have been reported during use of INVOKANA. Instruct patients to check ketones (when possible) if symptoms consistent with ketoacidosis occur even if blood glucose is not elevated. If symptoms of ketoacidosis (including nausea, vomiting, abdominal pain, tiredness, and labored breathing) occur, instruct patients to discontinue INVOKANA and seek medical advice immediately [see Warnings and Precautions (5.2)].

#### Acute Kidney Injury

Inform patients that acute kidney injury has been reported during use of INVOKANA. Advise patients to seek medical advice immediately if they have reduced oral intake (such as due to acute illness or fasting) or increased fluid losses (such as due to vomiting, diarrhea, or excessive heat exposure), as it may be appropriate to temporarily discontinue INVOKANA use in those settings [see Warnings and Precautions (5.3)].

**INVOKANA® (canagliflozin) tablets****Serious Urinary Tract Infections**

Inform patients of the potential for urinary tract infections, which may be serious. Provide them with information on the symptoms of urinary tract infections. Advise them to seek medical advice if such symptoms occur [see *Warnings and Precautions (5.5)*].

**Genital Mycotic Infections In Females (e.g., Vulvovaginitis)**

Inform female patients that vaginal yeast infection may occur and provide them with information on the signs and symptoms of vaginal yeast infection. Advise them of treatment options and when to seek medical advice [see *Warnings and Precautions (5.7)*].

**Genital Mycotic Infections In Males (e.g., Balanitis or Balanoposthitis)**

Inform male patients that yeast infection of penis (e.g., balanitis or balanoposthitis) may occur, especially in uncircumcised males and patients with prior history. Provide them with information on the signs and symptoms of balanitis and balanoposthitis (rash or redness of the glans or foreskin of the penis). Advise them of treatment options and when to seek medical advice [see *Warnings and Precautions (5.7)*].

**Hypersensitivity Reactions**

Inform patients that serious hypersensitivity reactions, such as urticaria, rash, anaphylaxis, and angioedema, have been reported with INVOKANA. Advise patients to report immediately any signs or symptoms suggesting allergic reaction, and to discontinue drug until they have consulted prescribing physicians.

**Bone Fracture**

Inform patients that bone fractures have been reported in patients taking INVOKANA. Provide them with information on factors that may contribute to fracture risk.

**Pregnancy**

Advise pregnant women, and females of reproductive potential of the potential risk to a fetus with treatment with INVOKANA [see *Use in Specific Populations (8.1)*]. Instruct females of reproductive potential to report pregnancies to their physicians as soon as possible.

**Lactation**


Advise women that breastfeeding is not recommended during treatment with INVOKANA [see *Use in Specific Populations (8.2)*].

Active ingredient made in Belgium

Manufactured for:  
Janssen Pharmaceuticals, Inc.  
Titusville, NJ 08560

Finished product manufactured by:  
Janssen Ortho, LLC  
Gurabo, PR 00778

Licensed from Mitsubishi Tanabe Pharma Corporation  
© 2013 Janssen Pharmaceuticals, Inc.



058051-160811

**Medication Guide**  
**INVOKANA® (In-vo-KAHN-uh)**  
**(canagliflozin)**  
**Tablets**

**What is the most important information I should know about INVOKANA?**

INVOKANA can cause important side effects, including:

- **Dehydration. INVOKANA can cause some people to become dehydrated (the loss of too much body water). Dehydration may cause you to feel dizzy, faint, lightheaded, or weak, especially when you stand up (orthostatic hypotension).**

You may be at higher risk of dehydration if you:

- have low blood pressure
- take medicines to lower your blood pressure, including diuretics (water pill)
- are on a low sodium (salt) diet
- have kidney problems
- are 65 years of age or older

Talk to your doctor about what you can do to prevent dehydration including how much fluid you should drink on a daily basis.

- **Vaginal yeast infection.** Women who take INVOKANA may get vaginal yeast infections. Symptoms of a vaginal yeast infection include:

- vaginal odor
- white or yellowish vaginal discharge (discharge may be lumpy or look like cottage cheese)
- vaginal itching

- **Yeast infection of the penis (balanitis or balanoposthitis).** Men who take INVOKANA may get a yeast infection of the skin around the penis. Certain men who are not circumcised may have swelling of the penis that makes it difficult to pull back the skin around the tip of the penis. Other symptoms of yeast infection of the penis include:

- redness, itching, or swelling of the penis
- rash of the penis
- foul smelling discharge from the penis
- pain in the skin around penis

Talk to your doctor about what to do if you get symptoms of a yeast infection of the vagina or penis. Your doctor may suggest you use an over-the-counter antifungal medicine. Talk to your doctor right away if you use an over-the-counter antifungal medication and your symptoms do not go away.

**What is INVOKANA?**

- INVOKANA is a prescription medicine used along with diet and exercise to lower blood sugar in adults with type 2 diabetes.
- INVOKANA is not for people with type 1 diabetes.
- INVOKANA is not for people with diabetic ketoacidosis (increased ketones in blood or urine).
- It is not known if INVOKANA is safe and effective in children under 18 years of age.

**Who should not take INVOKANA?**

**Do not take INVOKANA if you:**

- are allergic to canagliflozin or any of the ingredients in INVOKANA. See the end of this Medication Guide for a list of ingredients in INVOKANA. Symptoms of allergic reaction to INVOKANA may include:
  - rash
  - raised red patches on your skin (hives)
  - swelling of the face, lips, mouth, tongue, and throat that may cause difficulty in breathing or swallowing
- have severe kidney problems or are on dialysis.

**INVOKANA® (canagliflozin) tablets****What should I tell my doctor before taking INVOKANA?**

Before you take INVOKANA, tell your doctor if you:

- have kidney problems.
- have liver problems.
- have a history of urinary tract infections or problems with urination.
- are on a low sodium (salt) diet. Your doctor may change your diet or your dose of INVOKANA.
- are going to have surgery.
- are eating less due to illness, surgery, or a change in your diet.
- have or have had problems with your pancreas, including pancreatitis or surgery on your pancreas.
- drink alcohol very often, or drink a lot of alcohol in the short-term ("binge" drinking).
- have ever had an allergic reaction to INVOKANA.
- have other medical conditions.
- are pregnant or plan to become pregnant. INVOKANA may harm your unborn baby. If you become pregnant while taking INVOKANA, tell your doctor as soon as possible. Talk with your doctor about the best way to control your blood sugar while you are pregnant.
- are breastfeeding or plan to breastfeed. INVOKANA may pass into your breast milk and may harm your baby. Talk with your doctor about the best way to feed your baby if you are taking INVOKANA. Do not breastfeed while taking INVOKANA.

**Tell your doctor about all the medicines you take, including prescription and non-prescription medicines, vitamins, and herbal supplements.**

INVOKANA may affect the way other medicines work, and other medicines may affect how INVOKANA works. Especially tell your doctor if you take:

- diuretics (water pills)
- phenytoin or phenobarbital (used to control seizures)
- digoxin (Lanoxin®)\* (used to treat heart problems)
- rifampin (used to treat or prevent tuberculosis)
- ritonavir (Norvir®, Kaletra®)\* (used to treat HIV infection)

Ask your doctor or pharmacist for a list of these medicines if you are not sure if your medicine is listed above.

Know the medicines you take. Keep a list of them and show it to your doctor and pharmacist when you get a new medicine.

**How should I take INVOKANA?**

- Take INVOKANA by mouth 1 time each day exactly as your doctor tells you to take it.
- Your doctor will tell you how much INVOKANA to take and when to take it. Your doctor may change your dose if needed.
- It is best to take INVOKANA before the first meal of the day.
- Your doctor may tell you to take INVOKANA along with other diabetes medicines. Low blood sugar can happen more often when INVOKANA is taken with certain other diabetes medicines. See **"What are the possible side effects of INVOKANA?"**
- If you miss a dose, take it as soon as you remember. If it is almost time for your next dose, skip the missed dose and take the medicine at the next regularly scheduled time. Do not take two doses of INVOKANA at the same time. Talk to your doctor if you have questions about a missed dose.
- If you take too much INVOKANA, call your doctor or go to the nearest hospital emergency room right away.
- When your body is under some types of stress, such as fever, trauma (such as a car accident), infection, or surgery, the amount of diabetes medicine you need may change. Tell your doctor right away if you have any of these conditions and follow your doctor's instructions.
- Stay on your prescribed diet and exercise program while taking INVOKANA.
- Check your blood sugar as your doctor tells you to.
- INVOKANA will cause your urine to test positive for glucose.
- Your doctor may do certain blood tests before you start INVOKANA and during treatment as needed. Your doctor may change your dose of INVOKANA based on the results of your blood tests.
- Your doctor will check your diabetes with regular blood tests, including your blood sugar levels and your hemoglobin A1C.

**INVOKANA®** (canagliflozin) tablets**What are the possible side effects of INVOKANA?**

**INVOKANA may cause serious side effects including:**

See **"What is the most important information I should know about INVOKANA?"**

- **ketoacidosis (increased ketones in your blood or urine).** Ketoacidosis has happened in people who have **type 1 diabetes or type 2 diabetes**, during treatment with INVOKANA. Ketoacidosis is a serious condition, which may need to be treated in a hospital. Ketoacidosis may lead to death. **Ketoacidosis can happen with INVOKANA even if your blood sugar is less than 250 mg/dL. Stop taking INVOKANA and call your doctor right away if you get any of the following symptoms:**
  - nausea
  - vomiting
  - stomach area (abdominal) pain
  - tiredness
  - trouble breathing

If you get any of these symptoms during treatment with INVOKANA, if possible, check for ketones in your urine, even if your blood sugar is less than 250 mg/dL.

- **kidney problems.** Sudden kidney injury has happened to people taking INVOKANA. Talk to your doctor right away if you:
  - reduce the amount of food or liquid you drink for example, if you are sick or cannot eat or
  - you start to lose liquids from your body for example, from vomiting, diarrhea or being in the sun too long
- **a high amount of potassium in your blood (hyperkalemia)**
- **serious urinary tract infections.** Serious urinary tract infections that may lead to hospitalization have happened in people who are taking INVOKANA. Tell your doctor if you have any signs or symptoms of a urinary tract infection such as a burning feeling when passing urine, a need to urinate often, the need to urinate right away, pain in the lower part of your stomach (pelvis), or blood in the urine. Sometimes people may also have a fever, back pain, nausea, or vomiting.
- **low blood sugar (hypoglycemia).** If you take INVOKANA with another medicine that can cause low blood sugar, such as a sulfonylurea or insulin, your risk of getting low blood sugar is higher. The dose of your sulfonylurea medicine or insulin may need to be lowered while you take INVOKANA.

Signs and symptoms of low blood sugar may include:

- headache
- drowsiness
- weakness
- confusion
- dizziness
- irritability
- hunger
- fast heartbeat
- sweating
- shaking or feeling jittery
- **serious allergic reaction.** If you have any symptoms of a serious allergic reaction, stop taking INVOKANA and call your doctor right away or go to the nearest hospital emergency room. See **"Who should not take INVOKANA?"**. Your doctor may give you a medicine for your allergic reaction and prescribe a different medicine for your diabetes.
- **broken bones (fractures).** Bone fractures have been seen in patients taking INVOKANA. Talk to your doctor about factors that may increase your risk of bone fracture.

The most common side effects of INVOKANA include:

- vaginal yeast infections and yeast infections of the penis (See **"What is the most important information I should know about INVOKANA?"**)
- changes in urination, including urgent need to urinate more often, in larger amounts, or at night

Tell your doctor if you have any side effect that bothers you or that does not go away. These are not all the possible side effects of INVOKANA. For more information, ask your doctor or pharmacist.

**Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.**

You may also report side effects to Janssen Pharmaceuticals, Inc. at 1-800-526-7736.

**How should I store INVOKANA?**

- Store INVOKANA at room temperature between 68°F to 77°F (20°C to 25°C).
- **Keep INVOKANA and all medicines out of the reach of children.**

**INVOKANA® (canagliflozin) tablets****General information about the safe and effective use of INVOKANA.**

Medicines are sometimes prescribed for purposes other than those listed in the Medication Guide. Do not use INVOKANA for a condition for which it was not prescribed. Do not give INVOKANA to other people, even if they have the same symptoms you have. It may harm them.

This Medication Guide summarizes the most important information about INVOKANA. If you would like more information, talk with your doctor. You can ask your pharmacist or doctor for information about INVOKANA that is written for healthcare professionals.

**For more information about INVOKANA, call 1-800-526-7736 or visit our website at [www.invokana.com](http://www.invokana.com).**

**What are the ingredients of INVOKANA?**

Active ingredient: canagliflozin

Inactive ingredients: croscarmellose sodium, hydroxypropyl cellulose, lactose anhydrous, magnesium stearate, and microcrystalline cellulose. In addition, the tablet coating contains iron oxide yellow E172 (100 mg tablet only), macrogol/PEG, polyvinyl alcohol, talc, and titanium dioxide.

\* The brands listed are trademarks of their respective owners and are not trademarks of Janssen Pharmaceuticals, Inc. Active ingredient made in Belgium. Manufactured for: Janssen Pharmaceuticals, Inc., Titusville, NJ 08560. Manufactured by: Janssen Ortho, LLC, Gurabo, PR 00778. Licensed from Mitsubishi Tanabe Pharma Corporation. © 2013 Janssen Pharmaceuticals, Inc.

This Medication Guide has been approved by the U.S. Food and Drug Administration.

Revised 07/2016

058051-160811

**INVOKAMET®**  
(canagliflozin and metformin hydrochloride)  
tablets for oral use

Revised: 07/2016  
058065-160811

**HIGHLIGHTS OF PRESCRIBING INFORMATION**

These highlights do not include all the information needed to use INVOKAMET® safely and effectively. See full prescribing information for INVOKAMET.

INVOKAMET (canagliflozin and metformin hydrochloride) tablets, for oral use  
Initial U.S. Approval – 2014

**WARNING: LACTIC ACIDOSIS**

See full prescribing information for complete boxed warning.

- Postmarketing cases of metformin-associated lactic acidosis have resulted in death, hypothermia, hypotension, and resistant bradyarrhythmias. Symptoms included malaise, myalgias, respiratory distress, somnolence, and abdominal pain. Laboratory abnormalities included elevated blood lactate levels, anion gap acidosis, increased lactate/pyruvate ratio, and metformin plasma levels generally >5 mcg/mL. (5.1)
- Risk factors include renal impairment, concomitant use of certain drugs, age >65 years old, radiological studies with contrast, surgery and other procedures, hypoxic states, excessive alcohol intake, and hepatic impairment. Steps to reduce the risk of and manage metformin-associated lactic acidosis in these high risk groups are provided in the Full Prescribing Information. (5.1)
- If lactic acidosis is suspected, discontinue INVOKAMET and institute general supportive measures in a hospital setting. Prompt hemodialysis is recommended. (5.1)

**RECENT MAJOR CHANGES**

Boxed Warning	05/2016
Indications and Usage (1)	05/2016
Dosage and Administration (2)	05/2016
Contraindications (4)	05/2016
Warnings and Precautions (5)	07/2016

**INDICATIONS AND USAGE**

INVOKAMET is a sodium-glucose co-transporter 2 (SGLT2) inhibitor and biguanide combination product indicated as an adjunct to diet and exercise to improve glycemic control in adults with type 2 diabetes mellitus when treatment with both canagliflozin and metformin is appropriate (1)

**Limitation of use:**

Not for treatment of type 1 diabetes or diabetic ketoacidosis (1)

**DOSAGE AND ADMINISTRATION**

- Individualize based on the patient's current regimen (2)
- Take one INVOKAMET tablet twice daily with meals, recommended starting dose of canagliflozin is 50 mg twice daily and metformin 500 mg twice daily (2.1)
- Canagliflozin dose can be increased to 150 mg twice daily in patients tolerating canagliflozin 50 mg twice daily who have eGFR of 60 mL/min/1.73 m<sup>2</sup> or greater and require additional glycemic control. Do not exceed a total daily canagliflozin dose of 300 mg (2.1)
- Gradually escalate metformin dose to reduce the gastrointestinal side effects while not exceeding total daily dose of 2000 mg (2.1)
- Assess renal function before initiating and periodically thereafter (2.2)
- INVOKAMET is contraindicated in patients with an estimated glomerular filtration rate (eGFR) below 45 mL/min/1.73 m<sup>2</sup> (2.2)
- Limit the dose of canagliflozin component to 50 mg twice daily in patients with an eGFR of 45 to less than 60 mL/min/1.73 m<sup>2</sup> (2.2)
- INVOKAMET may need to be discontinued at time of, or prior to, iodinated contrast imaging procedures (2.4)

**DOSAGE FORMS AND STRENGTHS**

Film-coated tablets:

- Canagliflozin 50 mg and metformin hydrochloride 500 mg
- Canagliflozin 50 mg and metformin hydrochloride 1,000 mg
- Canagliflozin 150 mg and metformin hydrochloride 500 mg
- Canagliflozin 150 mg and metformin hydrochloride 1,000 mg (3)

**INVOKAMET®** (canagliflozin and metformin hydrochloride) tablets

**CONTRAINDICATIONS**

- Moderate to severe renal impairment (eGFR below 45 mL/min/1.73 m<sup>2</sup>), end stage renal disease or dialysis (4, 5.1, 5.4)
- Metabolic acidosis, including diabetic ketoacidosis (1, 4, 5.1)
- History of serious hypersensitivity reaction to canagliflozin or metformin (4, 5.8)

**WARNINGS AND PRECAUTIONS**

- **Lactic acidosis:** See boxed warning (5.1)
- **Hypotension:** Before initiating INVOKAMET, assess volume status and correct hypovolemia in patients with renal impairment, the elderly, in patients with low systolic blood pressure, or on diuretics, ACEi, or ARB. Monitor for signs and symptoms during therapy (5.2)
- **Ketoacidosis:** Assess patients who present with signs and symptoms of metabolic acidosis for ketoacidosis, regardless of blood glucose level. If suspected, discontinue INVOKAMET, evaluate and treat promptly. Before initiating INVOKAMET, consider risk factors for ketoacidosis. Patients on INVOKAMET may require monitoring and temporary discontinuation of therapy in clinical situations known to predispose to ketoacidosis (5.3)
- **Acute kidney injury and impairment in renal function:** Consider temporarily discontinuing in settings of reduced oral intake or fluid losses. If acute kidney injury occurs, discontinue and promptly treat. Monitor renal function during therapy (5.4)
- **Hyperkalemia:** Monitor potassium levels in patients with impaired renal function and in patients predisposed to hyperkalemia (2.2, 5.5, 6.1, 6.6)
- **Urosepsis and Pyelonephritis:** Evaluate patients for signs and symptoms of urinary tract infections and treat promptly, if indicated (5.6)
- **Hypoglycemia:** Consider a lower dose of insulin or the insulin secretagogue to reduce the risk of hypoglycemia when used in combination with INVOKAMET (5.7)
- **Genital mycotic infections:** Monitor and treat if indicated (5.8)
- **Hypersensitivity reactions:** Discontinue INVOKAMET and monitor until signs and symptoms resolve (5.9)
- **Bone fracture:** Consider factors that contribute to fracture risk before initiating INVOKAMET (5.10)
- **Vitamin B<sub>12</sub> deficiency:** Metformin may lower vitamin B<sub>12</sub> levels. Monitor hematologic parameters annually (5.11)
- **Increased LDL-C:** Monitor LDL-C and treat if appropriate (5.12)

**ADVERSE REACTIONS**

- Most common adverse reactions associated with canagliflozin (5% or greater incidence): female genital mycotic infections, urinary tract infection, and increased urination (6.1)
- Most common adverse reactions associated with metformin (5% or greater incidence) are diarrhea, nausea, vomiting, flatulence, asthenia, indigestion, abdominal discomfort, and headache (6.1)

To report SUSPECTED ADVERSE REACTIONS, contact Janssen Pharmaceuticals, Inc. at 1-800-526-7736 or FDA at 1-800-FDA-1088 or [www.fda.gov/medwatch](http://www.fda.gov/medwatch).

**DRUG INTERACTIONS**

- **Carbonic anhydrase inhibitors** may increase risk of lactic acidosis. Consider more frequent monitoring (7.1)
- **Drugs that are eliminated by renal tubular secretion** (e.g. cationic drugs such as cimetidine), may increase the accumulation of metformin. Consider more frequent monitoring (7.1)
- **Alcohol** can potentiate the effect of metformin on lactate metabolism. Warn patients against excessive alcohol intake (7.1)
- **UGT inducers** (e.g., rifampin): Canagliflozin exposure is reduced. Consider increasing canagliflozin dose from 50 mg to 150 mg twice daily (2.3, 7.2)
- **Digoxin:** Monitor digoxin levels (7.2)

**USE IN SPECIFIC POPULATIONS**

- **Pregnancy:** Advise females of the potential risk to a fetus especially during the second and third trimesters (8.1)
- **Lactation:** INVOKAMET is not recommended when breastfeeding (8.2)
- **Females and Males of Reproductive Potential:** Advise premenopausal females of the potential for an unintended pregnancy (8.3)
- **Geriatrics:** Higher incidence of adverse reactions related to reduced intravascular volume. Assess renal function more frequently (5.2, 6.1, 8.5)
- **Renal impairment:** Higher incidence of adverse reactions related to reduced intravascular volume and renal function (2.2, 5.4, 8.6)
- **Hepatic impairment:** Avoid use in patients with hepatic impairment (8.7)

See 17 for PATIENT COUNSELING INFORMATION and Medication Guide.

Revised: 07/2016

**INVOKAMET® (canagliflozin and metformin hydrochloride) tablets****INVOKAMET® (canagliflozin and metformin hydrochloride) tablets****FULL PRESCRIBING INFORMATION: CONTENTS\*****WARNING: LACTIC ACIDOSIS****1 INDICATIONS AND USAGE****2 DOSAGE AND ADMINISTRATION**

- 2.1 Recommended Dosage
- 2.2 Recommended Dosage for Patients with Renal Impairment
- 2.3 Concomitant Use with UDP-Glucuronosyl Transferase (UGT) Enzyme Inducers
- 2.4 Discontinuation for Iodinated Contrast Imaging Procedures

**3 DOSAGE FORMS AND STRENGTHS****4 CONTRAINDICATIONS****5 WARNINGS AND PRECAUTIONS**

- 5.1 Lactic Acidosis
- 5.2 Hypotension
- 5.3 Ketoacidosis
- 5.4 Acute Kidney Injury and Impairment in Renal Function
- 5.5 Hyperkalemia
- 5.6 Urosepsis and Pyelonephritis
- 5.7 Hypoglycemia with Concomitant Use of Sulfonylurea or Insulin
- 5.8 Genital Mycotic Infections
- 5.9 Hypersensitivity Reactions
- 5.10 Bone Fracture
- 5.11 Vitamin B<sub>12</sub> Levels
- 5.12 Increases in Low-Density Lipoprotein (LDL-C)
- 5.13 Macrovascular Outcomes

**6 ADVERSE REACTIONS**

- 6.1 Clinical Studies Experience
- 6.2 Postmarketing Experience

**7 DRUG INTERACTIONS**

- 7.1 Drug Interactions with Metformin
- 7.2 Drug Interactions with Canagliflozin

**8 USE IN SPECIFIC POPULATIONS**

- 8.1 Pregnancy
- 8.2 Lactation
- 8.3 Females and Males of Reproductive Potential
- 8.4 Pediatric Use
- 8.5 Geriatric Use
- 8.6 Renal Impairment
- 8.7 Hepatic Impairment

**10 OVERDOSAGE****11 DESCRIPTION****12 CLINICAL PHARMACOLOGY**

- 12.1 Mechanism of Action
- 12.2 Pharmacodynamics
- 12.3 Pharmacokinetics

**13 NONCLINICAL TOXICOLOGY**

- 13.1 Carcinogenesis, Mutagenesis, Impairment of Fertility

**14 CLINICAL STUDIES**

- 14.1 Canagliflozin as Initial Combination Therapy with Metformin
- 14.2 Canagliflozin as Add-on Combination Therapy with Metformin
- 14.3 Canagliflozin Compared to Glimepiride, Both as Add-on Combination Therapy with Metformin
- 14.4 Canagliflozin as Add-on Combination Therapy with Metformin and Sulfonylurea
- 14.5 Canagliflozin Compared to Sitagliptin, Both as Add-on Combination Therapy with Metformin and Sulfonylurea
- 14.6 Canagliflozin as Add-on Combination Therapy with Metformin and Pioglitazone
- 14.7 Canagliflozin as Add-on Combination Therapy with Insulin (With or Without Other Anti-Hyperglycemic Agents, Including Metformin)

**16 HOW SUPPLIED/STORAGE AND HANDLING****17 PATIENT COUNSELING INFORMATION**

\*Sections or subsections omitted from the full prescribing information are not listed.

**FULL PRESCRIBING INFORMATION****WARNING: LACTIC ACIDOSIS**

- Post-marketing cases of metformin-associated lactic acidosis have resulted in death, hypothermia, hypotension, and resistant bradyarrhythmias. The onset of metformin-associated lactic acidosis is often subtle, accompanied only by nonspecific symptoms such as malaise, myalgias, respiratory distress, somnolence, and abdominal pain. Metformin-associated lactic acidosis was characterized by elevated blood lactate levels (> 5 mmol/Liter), anion gap acidosis (without evidence of ketonuria or ketonemia), an increased lactate/pyruvate ratio; and metformin plasma levels generally >5 mcg/mL. [see Warnings and Precautions (5.1)].
- Risk factors for metformin-associated lactic acidosis include renal impairment, concomitant use of certain drugs (e.g., cationic drugs such as topiramate), age 65 years old or greater, having a radiological study with contrast, surgery and other procedures, hypoxic states (e.g., acute congestive heart failure), excessive alcohol intake, and hepatic impairment.
- Steps to reduce the risk of and manage metformin-associated lactic acidosis in these high risk groups are provided in the full prescribing information [see Dosage and Administration (2.2), Contraindications (4), Warnings and Precautions (5.1), Drug Interactions (7), and Use in Specific Populations (8.6, 8.7)].
- If metformin-associated lactic acidosis is suspected, immediately discontinue INVOKAMET and institute general supportive measures in a hospital setting. Prompt hemodialysis is recommended [see Warnings and Precautions (5.1)].

**1 INDICATIONS AND USAGE**

INVOKAMET (canagliflozin and metformin hydrochloride) is indicated as an adjunct to diet and exercise to improve glycemic control in adults with type 2 diabetes mellitus when treatment with both canagliflozin and metformin is appropriate.

**Limitations of Use**

INVOKAMET is not recommended in patients with type 1 diabetes or for the treatment of diabetic ketoacidosis.

**2 DOSAGE AND ADMINISTRATION****2.1 Recommended Dosage**

- Individualize the starting dose of INVOKAMET (canagliflozin and metformin hydrochloride) based on the patient's current regimen:
  - In patients currently not treated with either canagliflozin or metformin, initiate therapy with INVOKAMET containing canagliflozin 50 mg and metformin 500 mg [see Clinical Studies (14.1)].

- In patients on metformin, switch to INVOKAMET containing canagliflozin 50 mg and the same, or nearest appropriate, daily dose of metformin;
- In patients on canagliflozin, switch to INVOKAMET containing metformin 500 mg with the same daily dose of canagliflozin;
- In patients already treated with canagliflozin and metformin, switch to INVOKAMET containing the same daily dose of canagliflozin and the same, or nearest appropriate, daily dose of metformin.

- Take one INVOKAMET tablet twice daily with meals; in patients tolerating canagliflozin 50 mg twice daily who have an eGFR of 60 mL/min/1.73 m<sup>2</sup> or greater and require additional glycemic control, INVOKAMET dose can be increased for the canagliflozin component to 150 mg twice daily, with gradual metformin dose escalation to reduce the gastrointestinal side effects due to metformin [see Dosage Forms and Strengths (3), Clinical Studies (14.1)].

- In patients with volume depletion not previously treated with canagliflozin, correct this condition before initiating INVOKAMET [see Warnings and Precautions (5.2), Use in Specific Populations (8.5, 8.6), and Patient Counseling Information (17)].

- Adjust dosing based on effectiveness and tolerability while not exceeding the maximum recommended daily dose of metformin 2000 mg and canagliflozin 300 mg in patients with an eGFR of 60 mL/min/1.73 m<sup>2</sup> or greater [see Dosage and Administration (2.2)].

**2.2 Recommended Dosage for Patients with Renal Impairment**

- Assess renal function before initiating INVOKAMET and periodically thereafter.
- INVOKAMET is contraindicated in patients with an estimated glomerular filtration rate (eGFR) below 45 mL/min/1.73 m<sup>2</sup> [see Contraindications (4), Warnings and Precautions (5.1, 5.3)].
- Limit the dose of the canagliflozin component to 50 mg twice daily in patients with moderate renal impairment with an eGFR of 45 to less than 60 mL/min/1.73 m<sup>2</sup>.

**2.3 Concomitant Use with UDP-Glucuronosyl Transferase (UGT) Enzyme Inducers**

If an inducer of UGTs (e.g., rifampin, phenytoin, phenobarbital, ritonavir) is co-administered with INVOKAMET, consider increasing the dose to canagliflozin 150 mg twice daily in patients currently tolerating canagliflozin 50 mg twice daily who have an eGFR of 60 mL/min/1.73 m<sup>2</sup> or greater and require additional glycemic control [see Drug Interactions (7.2)].

Consider another antihyperglycemic agent in patients with an eGFR of 45 to less than 60 mL/min/1.73 m<sup>2</sup> receiving concurrent therapy with a UGT inducer.

**INVOKAMET® (canagliflozin and metformin hydrochloride) tablets****2.4 Discontinuation for Iodinated Contrast Imaging Procedures**

Discontinue INVOKAMET at the time of, or prior to, an iodinated contrast imaging procedure in patients with an eGFR between 45 and 60 mL/min/1.73 m<sup>2</sup>. In patients with a history of liver disease, alcoholism or heart failure; or in patients who will be administered intra-arterial iodinated contrast. Re-evaluate eGFR 48 hours after the imaging procedure; restart INVOKAMET if renal function is stable [see *Warnings and Precautions* (5.1)].

**3 DOSAGE FORMS AND STRENGTHS**

INVOKAMET (canagliflozin and metformin hydrochloride) film-coated tablets for oral administration are available in the following strengths:

- Canagliflozin 50 mg and metformin hydrochloride 500 mg tablets are immediate-release, capsule-shaped, white film-coated tablets with "CM" on one side and "155" on the other side.
- Canagliflozin 50 mg and metformin hydrochloride 1,000 mg tablets are immediate-release, capsule-shaped, beige, film-coated tablets with "CM" on one side and "551" on the other side.
- Canagliflozin 150 mg and metformin hydrochloride 500 mg tablets are immediate-release, capsule-shaped, yellow, film-coated tablets with "CM" on one side and "215" on the other side.
- Canagliflozin 150 mg and metformin hydrochloride 1,000 mg tablets are immediate-release, capsule-shaped, purple, film-coated tablets with "CM" on one side and "611" on the other side.

**4 CONTRAINDICATIONS**

INVOKAMET is contraindicated in patients with:

- Moderate to severe renal impairment (eGFR below 45 mL/min/1.73 m<sup>2</sup>), end stage renal disease (ESRD) or patients on dialysis [see *Warnings and Precautions* (5.1) and *Use in Specific Populations* (8.6)].
- Acute or chronic metabolic acidosis, including diabetic ketoacidosis [see *Warnings and Precautions* (5.3)].
- History of a serious hypersensitivity reaction to canagliflozin or metformin, such as anaphylaxis or angioedema [see *Warnings and Precautions* (5.9) and *Adverse Reactions* (6.1, 6.2)].

**5 WARNINGS AND PRECAUTIONS****5.1 Lactic Acidosis**

There have been post-marketing cases of metformin-associated lactic acidosis, including fatal cases. These cases had a subtle onset and were accompanied by nonspecific symptoms such as malaise, myalgias, abdominal pain, respiratory distress, or increased somnolence; however, hypothermia, hypotension and resistant bradyarrhythmias have occurred with severe acidosis. Metformin-associated lactic acidosis was characterized by elevated blood lactate concentrations (>5 mmol/Liter), anion gap acidosis (without evidence of ketonuria or ketonemia), and an increased lactate:pyruvate ratio; metformin plasma levels generally >5 mcg/mL. Metformin decreases liver uptake of lactate increasing lactate blood levels which may increase the risk of lactic acidosis, especially in patients at risk.

If metformin-associated lactic acidosis is suspected, general supportive measures should be instituted promptly in a hospital setting, along with immediate discontinuation of INVOKAMET. In INVOKAMET-treated patients with a diagnosis or strong suspicion of lactic acidosis, prompt hemodialysis is recommended to correct the acidosis and remove accumulated metformin (metformin hydrochloride is dialyzable, with a clearance of up to 170 mL/minute under good hemodynamic conditions). Hemodialysis has often resulted in reversal of symptoms and recovery.

Educate patients and their families about the symptoms of lactic acidosis and if these symptoms occur instruct them to discontinue INVOKAMET and report these symptoms to their healthcare provider.

For each of the known and possible risk factors for metformin-associated lactic acidosis, recommendations to reduce the risk of and manage metformin-associated lactic acidosis are provided below:

**Renal Impairment:** The postmarketing metformin-associated lactic acidosis cases primarily occurred in patients with significant renal impairment. The risk of metformin accumulation and metformin-associated lactic acidosis increases with the severity of renal impairment because metformin is substantially excreted by the kidney [see *Clinical Pharmacology* (12.3)].

- Before initiating INVOKAMET, obtain an estimated glomerular filtration rate (eGFR).
- INVOKAMET is contraindicated in patients with an eGFR less than 45 mL/minute/1.73 m<sup>2</sup>.
- Obtain an eGFR at least annually in all patients taking INVOKAMET. In patients at increased risk for the development of renal impairment (e.g., the elderly), renal function should be assessed more frequently.

**Drug Interactions:** The concomitant use of INVOKAMET with specific drugs may increase the risk of metformin-associated lactic acidosis; those that impair renal function, result in significant hemodynamic change, interfere with acid-base balance or increase metformin accumulation (e.g. cationic drugs) [see *Drug Interactions* (7)]. Therefore, consider more frequent monitoring of patients.

**INVOKAMET® (canagliflozin and metformin hydrochloride) tablets**

**Age 65 or Greater:** The risk of metformin-associated lactic acidosis increases with the patient's age because elderly patients have a greater likelihood of having hepatic, renal, or cardiac impairment than younger patients. Assess renal function more frequently in elderly patients [see *Use in Specific Populations* (8.5)].

**Radiological Studies with Contrast:** Administration of intravascular iodinated contrast agents in metformin-treated patients has led to an acute decrease in renal function and the occurrence of lactic acidosis. Stop INVOKAMET at the time of, or prior to, an iodinated contrast imaging procedure in patients with an eGFR between 45 and 60 mL/min/1.73 m<sup>2</sup>. In patients with a history of hepatic impairment, alcoholism, or heart failure; or in patients who will be administered intra-arterial iodinated contrast. Re-evaluate eGFR 48 hours after the imaging procedure, and restart INVOKAMET if renal function is stable.

**Surgery and Other Procedures:** Withholding of food and fluids during surgical or other procedures may increase the risk for volume depletion, hypotension and renal impairment.

INVOKAMET should be temporarily discontinued while patients have restricted food and fluid intake.

**Hypoxic States:** Several of the postmarketing cases of metformin-associated lactic acidosis occurred in the setting of acute congestive heart failure (particularly when accompanied by hypoperfusion and hypoxemia). Cardiovascular collapse (shock), acute myocardial infarction, sepsis, and other conditions associated with hypoxemia have been associated with lactic acidosis and may also cause prerenal azotemia. When such events occur, discontinue INVOKAMET.

**Excessive Alcohol Intake:** Alcohol potentiates the effect of metformin on lactate metabolism and this may increase the risk of metformin-associated lactic acidosis. Warn patients against excessive alcohol intake while receiving INVOKAMET.

**Hepatic Impairment:** Patients with hepatic impairment have developed metformin-associated lactic acidosis. This may be due to impaired lactate clearance resulting in higher lactate blood levels. Therefore, avoid use of INVOKAMET in patients with clinical or laboratory evidence of hepatic disease.

**5.2 Hypotension**

Canagliflozin causes intravascular volume contraction. Symptomatic hypotension can occur after initiating INVOKAMET [see *Adverse Reactions* (6.1)] particularly in patients with eGFR less than 60 mL/min/1.73 m<sup>2</sup>, elderly patients, patients on either diuretics or medications that interfere with the renin-angiotensin-aldosterone system (e.g., angiotensin-converting-enzyme [ACE] inhibitors, angiotensin receptor blockers [ARBs]), or patients with low systolic blood pressure. Before initiating INVOKAMET in patients with one or more of these characteristics who were not already on canagliflozin, volume status should be assessed and corrected. Monitor for signs and symptoms after initiating therapy.

**5.3 Ketoacidosis**

Reports of ketoacidosis, a serious life-threatening condition requiring urgent hospitalization have been identified in postmarketing surveillance in patients with type 1 and type 2 diabetes mellitus receiving sodium glucose co-transporter-2 (SGLT2) inhibitors, including canagliflozin. Fatal cases of ketoacidosis have been reported in patients taking canagliflozin. INVOKAMET is not indicated for the treatment of patients with type 1 diabetes mellitus [see *Indications and Usage* (1)].

Patients treated with INVOKAMET who present with signs and symptoms consistent with severe metabolic acidosis should be assessed for ketoacidosis regardless of presenting blood glucose levels, as ketoacidosis associated with INVOKAMET may be present even if blood glucose levels are less than 250 mg/dL. If ketoacidosis is suspected, INVOKAMET should be discontinued, patient should be evaluated, and prompt treatment should be instituted. Treatment of ketoacidosis may require insulin, fluid and carbohydrate replacement.

In many of the postmarketing reports, and particularly in patients with type 1 diabetes, the presence of ketoacidosis was not immediately recognized and institution of treatment was delayed because presenting blood glucose levels were below those typically expected for diabetic ketoacidosis (often less than 250 mg/dL). Signs and symptoms at presentation were consistent with dehydration and severe metabolic acidosis and included nausea, vomiting, abdominal pain, generalized malaise, and shortness of breath. In some but not all cases, factors predisposing to ketoacidosis such as insulin dose reduction, acute febrile illness, reduced caloric intake due to illness or surgery, pancreatic disorders suggesting insulin deficiency (e.g., type 1 diabetes, history of pancreatitis or pancreatic surgery), and alcohol abuse were identified.

Before initiating INVOKAMET consider factors in the patient history that may predispose to ketoacidosis including pancreatic insulin deficiency from any cause, caloric restriction, and alcohol abuse. In patients treated with INVOKAMET consider monitoring for ketoacidosis and temporarily discontinuing INVOKAMET in clinical situations known to predispose to ketoacidosis (e.g., prolonged fasting due to acute illness or surgery).

**5.4 Acute Kidney Injury and Impairment in Renal Function**

Canagliflozin causes intravascular volume contraction [see *Warnings and Precautions* (5.2)] and can cause renal impairment [see *Adverse Reactions* (6.1)]. There have been postmarketing reports of acute kidney injury, some requiring hospitalization and dialysis, in patients receiving canagliflozin; some reports involved patients younger than 65 years of age.

**INVOKAMET® (canagliflozin and metformin hydrochloride) tablets**

Before initiating INVOKAMET, consider factors that may predispose patients to acute kidney injury including hypovolemia, chronic renal insufficiency, congestive heart failure, and concomitant medications (diuretics, ACE inhibitors, ARBs, NSAIDs). Consider temporarily discontinuing INVOKAMET in any setting of reduced oral intake (such as acute illness or fasting) or fluid losses (such as gastrointestinal illness or excessive heat exposure); monitor patients for signs and symptoms of acute kidney injury. If acute kidney injury occurs, discontinue INVOKAMET promptly and institute treatment.

Canagliflozin increases serum creatinine and decreases eGFR. Patients with hypovolemia may be more susceptible to these changes. Renal function abnormalities can occur after initiating INVOKAMET [see *Adverse Reactions* (6.1)]. Renal function should be evaluated prior to initiation of INVOKAMET and monitored periodically thereafter. Dosage adjustment and more frequent renal function monitoring are recommended in patients with an eGFR below 60 mL/min/1.73 m<sup>2</sup>. INVOKAMET is contraindicated in patients with an eGFR below 45 mL/min/1.73 m<sup>2</sup> [see *Dosage and Administration* (2.2), *Contraindications* (4), *Warnings and Precautions* (5.1), *Use in Specific Populations* (8.6)].

**5.5 Hyperkalemia**

Canagliflozin can lead to hyperkalemia. Patients with moderate renal impairment who are taking medications that interfere with potassium excretion, such as potassium-sparing diuretics, or medications that interfere with the renin-angiotensin-aldosterone system are at an increased risk of developing hyperkalemia [see *Dosage and Administration* (2.2) and *Adverse Reactions* (6.1)].

Monitor serum potassium levels periodically after initiating INVOKAMET in patients with impaired renal function and in patients predisposed to hyperkalemia due to medications or other medical conditions.

**5.6 Urosepsis and Pyelonephritis**

There have been postmarketing reports of serious urinary tract infections including urosepsis and pyelonephritis requiring hospitalization in patients receiving SGLT2 inhibitors, including canagliflozin. Treatment with SGLT2 inhibitors increases the risk for urinary tract infections. Evaluate patients for signs and symptoms of urinary tract infections and treat promptly, if indicated [see *Adverse Reactions* (6)].

**5.7 Hypoglycemia with Concomitant Use of Sulfonylurea or Insulin****Canagliflozin**

Insulin and insulin secretagogues are known to cause hypoglycemia. Canagliflozin can increase the risk of hypoglycemia when combined with insulin or an insulin secretagogue [see *Adverse Reactions* (6.1)]. Therefore, a lower dose of insulin or insulin secretagogue may be required to minimize the risk of hypoglycemia when used in combination with INVOKAMET.

**Metformin**

Hypoglycemia does not occur in patients receiving metformin alone under usual circumstances of use, but could occur when caloric intake is deficient, when strenuous exercise is not compensated by caloric supplementation, or during concomitant use with other glucose-lowering agents (such as sulfonylureas and insulin) or ethanol. Elderly, debilitated, or malnourished patients, and those with adrenal or pituitary insufficiency or alcohol intoxication, are particularly susceptible to hypoglycemic effects. Hypoglycemia may be difficult to recognize in the elderly, and in people who are taking beta-adrenergic blocking drugs. Monitor for a need to lower the dose of INVOKAMET to minimize the risk of hypoglycemia in these patients.

**5.8 Genital Mycotic Infections**

Canagliflozin increases the risk of genital mycotic infections. Patients with a history of genital mycotic infections and uncircumcised males were more likely to develop genital mycotic infections [see *Adverse Reactions* (6.1)]. Monitor and treat appropriately.

**5.9 Hypersensitivity Reactions**

Hypersensitivity reactions, including angioedema and anaphylaxis, have been reported with canagliflozin. These reactions generally occurred within hours to days after initiating canagliflozin. If hypersensitivity reactions occur, discontinue use of INVOKAMET; treat and monitor until signs and symptoms resolve [see *Contraindications* (4) and *Adverse Reactions* (6.1, 6.2)].

**5.10 Bone Fracture**

An increased risk of bone fracture, occurring as early as 12 weeks after treatment initiation, was observed in patients using canagliflozin. Consider factors that contribute to fracture risk prior to initiating INVOKAMET [see *Adverse Reactions* (6.1)].

**INVOKAMET® (canagliflozin and metformin hydrochloride) tablets****5.11 Vitamin B<sub>12</sub> Levels**

In controlled, 29-week clinical trials of metformin, a decrease to subnormal levels of previously normal serum vitamin B<sub>12</sub> levels, without clinical manifestations, was observed in approximately 7% of metformin-treated patients. Such decreases, possibly due to interference with B<sub>12</sub> absorption from the B<sub>12</sub>-intrinsic factor complex, is, however, very rarely associated with anemia or neurologic manifestations due to the short duration (less than 1 year) of the clinical trials. This risk may be more relevant to patients receiving long-term treatment with metformin and adverse hematologic and neurologic reactions have been reported postmarketing. The decrease in vitamin B<sub>12</sub> levels appears to be rapidly reversible with discontinuation of metformin or vitamin B<sub>12</sub> supplementation. Measure hematologic parameters on an annual basis in patients on INVOKAMET and investigate and treat if abnormalities occur. Patients with inadequate vitamin B<sub>12</sub> or calcium intake or absorption may be predisposed to developing subnormal vitamin B<sub>12</sub> levels, and routine serum vitamin B<sub>12</sub> measurement at 2- to 3-year intervals is recommended in these patients.

**5.12 Increases in Low-Density Lipoprotein (LDL-C)**

Dose-related increases in LDL-C occur with canagliflozin [see *Adverse Reactions* (6.1)]. Monitor LDL-C and treat if appropriate after initiating INVOKAMET.

**5.13 Macrovascular Outcomes**

There have been no clinical studies establishing conclusive evidence of macrovascular risk reduction with INVOKAMET or any other antidiabetic drug [see *Adverse Reactions* (6.1)].

**6 ADVERSE REACTIONS**

The following adverse reactions are also discussed elsewhere in the labeling:

- Lactic Acidosis [see *Boxed Warning and Warnings and Precautions* (5.1, 5.4)]
- Hypotension [see *Warnings and Precautions* (5.2)]
- Ketoacidosis [see *Warnings and Precautions* (5.3)]
- Acute Kidney Injury and Impairment in Renal Function [see *Warnings and Precautions* (5.4)]
- Hyperkalemia [see *Warnings and Precautions* (5.5)]
- Urosepsis and Pyelonephritis [see *Warnings and Precautions* (5.6)]
- Hypoglycemia with Concomitant Use of Sulfonylurea or Insulin [see *Warnings and Precautions* (5.7)]
- Genital Mycotic Infections [see *Warnings and Precautions* (5.8)]
- Hypersensitivity Reactions [see *Warnings and Precautions* (5.9)]
- Bone Fracture [see *Warnings and Precautions* (5.10)]
- Vitamin B<sub>12</sub> Deficiency [see *Warnings and Precautions* (5.11)]
- Increases in Low-Density Lipoprotein (LDL-C) [see *Warnings and Precautions* (5.12)]

**6.1 Clinical Studies Experience**

Because clinical trials are conducted under widely varying conditions, adverse reaction rates observed in the clinical trials of a drug cannot be directly compared to the rates in the clinical trials of another drug and may not reflect the rates observed in clinical practice.

**Pool of Placebo-Controlled Trials****Canagliflozin**

The data in Table 1 is derived from four 28-week placebo-controlled trials. In one trial canagliflozin was used as monotherapy and in three trials canagliflozin was used as add-on therapy with metformin (with or without other agents) [see *Clinical Studies* (14)]. These data reflect exposure of 1667 patients to canagliflozin and a mean duration of exposure to canagliflozin of 24 weeks with 1275 patients exposed to a combination of canagliflozin and metformin. Patients received canagliflozin 100 mg (N=833), canagliflozin 300 mg (N=834) or placebo (N=846) once daily. The mean daily dose of metformin was 2136 mg (SD 337.3) for the 1275 patients in the three placebo-controlled metformin add-on studies. The mean age of the population was 56 years and 2% were older than 75 years of age. Fifty percent (50%) of the population was male and 72% were Caucasian, 12% were Asian, and 5% were Black or African American. At baseline the population had diabetes for an average of 7.3 years, had a mean HbA<sub>1c</sub> of 8.0% and 20% had established microvascular complications of diabetes. Baseline renal function was normal or mildly impaired (mean eGFR 88 mL/min/1.73 m<sup>2</sup>).

Table 1 shows common adverse reactions associated with the use of canagliflozin. These adverse reactions were not present at baseline, occurred more commonly on canagliflozin than on placebo, and occurred in at least 2% of patients treated with either canagliflozin 100 mg or canagliflozin 300 mg.

**INVOKAMET® (canagliflozin and metformin hydrochloride) tablets****Table 1: Adverse Reactions From Pool of Four 26-Week Placebo-Controlled Studies Reported in ≥ 2% of Canagliflozin-Treated Patients\***

Adverse Reaction	Placebo N=646	Canagliflozin 100 mg N=833	Canagliflozin 300 mg N=834
Urinary tract infections†	3.8%	5.9%	4.4%
Increased urination‡	0.7%	5.1%	4.6%
Thirst‡	0.1%	2.8%	2.4%
Constipation	0.9%	1.8%	2.4%
Nausea	1.6%	2.1%	2.3%
	N=312	N=425	N=430
Female genital mycotic infections†	2.8%	10.6%	11.6%
Vulvovaginal pruritus	0.0%	1.6%	3.2%
	N=334	N=409	N=404
Male genital mycotic infections†	0.7%	4.2%	3.8%

\* The four placebo-controlled trials included one monotherapy trial and three add-on combination trials with metformin, metformin and sulfonylurea, or metformin and pioglitazone.

† Female genital mycotic infections include the following adverse reactions: Vulvovaginal candidiasis, Vulvovaginal mycotic infection, Vulvovaginitis, Vaginal infection, Vulvitis, and Genital infection fungal.

‡ Urinary tract infections include the following adverse reactions: Urinary tract infection, Cystitis, Kidney infection, and Urosepsis.

§ Increased urination includes the following adverse reactions: Polyuria, Pollakiuria, Urine output increased, Micturition urgency, and Nocturia.

¶ Male genital mycotic infections include the following adverse reactions: Balanitis or Balanoposthitis, Balanitis candida, and Genital Infection fungal.

‡ Thirst includes the following adverse reactions: Thirst, Dry mouth, and Polydipsia. Note: Percentages were weighted by studies. Study weights were proportional to the harmonic mean of the three treatment sample sizes.

Abdominal pain was also more commonly reported in patients taking canagliflozin 100 mg (1.8%), 300 mg (1.7%) than in patients taking placebo (0.9%).

**Canagliflozin and Metformin**

The incidence and type of adverse reactions in the three 26-week placebo-controlled metformin add-on studies, representing a majority of data from the four 26-week placebo-controlled trials, was similar to the adverse reactions described in Table 1. There were no additional adverse reactions identified in the pooling of these three placebo-controlled studies that included metformin relative to the four placebo-controlled studies.

In a trial with canagliflozin as initial combination therapy with metformin [see *Clinical Studies* (14.1)], an increased incidence of diarrhea was observed in the canagliflozin and metformin combination groups (4.2%) compared to canagliflozin or metformin monotherapy groups (1.7%).

**Pool of Placebo- and Active-Controlled Trials - Canagliflozin**

The occurrence of adverse reactions for canagliflozin was evaluated in a larger pool of patients participating in placebo- and active-controlled trials.

The data combined eight clinical trials and reflect exposure of 8177 patients to canagliflozin. The mean duration of exposure to canagliflozin was 38 weeks with 1832 individuals exposed to canagliflozin for greater than 50 weeks. Patients received canagliflozin 100 mg (N=3092), canagliflozin 300 mg (N=3085) or comparator (N=3262) once daily. The mean age of the population was 60 years and 5% were older than 75 years of age. Fifty-eight percent (58%) of the population was male and 73% were Caucasian, 16% were Asian, and 4% were Black or African American. At baseline, the population had diabetes for an average of 11 years, had a mean HbA1C of 8.0% and 33% had established microvascular complications of diabetes. Baseline renal function was normal or mildly impaired (mean eGFR 81 mL/min/1.73 m<sup>2</sup>).

The types and frequency of common adverse reactions observed in the pool of eight clinical trials were consistent with those listed in Table 1. Percentages were weighted by studies. Study weights were proportional to the harmonic mean of the three treatment sample sizes. In this pool, canagliflozin was also associated with the adverse reactions of fatigue (1.8% with comparator, 2.2% with canagliflozin 100 mg, and 2.0% with canagliflozin 300 mg) and loss of strength or energy (i.e., asthenia) (0.6% with comparator, 0.7% with canagliflozin 100 mg, and 1.1% with canagliflozin 300 mg).

In the pool of eight clinical trials, the incidence rate of pancreatitis (acute or chronic) was 0.1, 0.2, and 0.1 receiving comparator, canagliflozin 100 mg, and canagliflozin 300 mg, respectively.

In the pool of eight clinical trials, hypersensitivity-related adverse reactions (including erythema, rash, pruritus, urticaria, and angioedema) occurred in 3.0%, 3.8%, and 4.2% of patients receiving comparator, canagliflozin 100 mg, and canagliflozin 300 mg, respectively. Five patients experienced serious adverse reactions of hypersensitivity with canagliflozin, which included 4 patients with urticaria and 1 patient with a diffuse rash and urticaria occurring within hours of exposure to canagliflozin. Among these patients, 2 patients discontinued canagliflozin. One patient with urticaria had recurrence when canagliflozin was re-initiated.

**INVOKAMET® (canagliflozin and metformin hydrochloride) tablets**

Photosensitivity-related adverse reactions (including photosensitivity reaction, polymorphic light eruption, and sunburn) occurred in 0.1%, 0.2%, and 0.2% of patients receiving comparator, canagliflozin 100 mg, and canagliflozin 300 mg, respectively.

Other adverse reactions occurring more frequently on canagliflozin than on comparator were:

**Volume Depletion-Related Adverse Reactions**

Canagliflozin results in an osmotic diuresis, which may lead to reductions in intravascular volume. In clinical studies, treatment with canagliflozin was associated with a dose-dependent increase in the incidence of volume depletion-related adverse reactions (e.g., hypotension, postural dizziness, orthostatic hypotension, syncope, and dehydration). An increased incidence was observed in patients on the 300 mg dose. The three factors associated with the largest increase in volume depletion-related adverse reactions were the use of loop diuretics, moderate renal impairment (eGFR 30 to less than 60 mL/min/1.73 m<sup>2</sup>), and age 75 years and older (Table 2) [see *Dosage and Administration* (2.2), *Warnings and Precautions* (5.2), and *Use in Specific Populations* (8.5, 8.6)].

**Table 2: Proportion of Patients With at Least One Volume Depletion-Related Adverse Reaction (Pooled Results from 8 Clinical Trials)**

Baseline Characteristic	Comparator Group* %	Canagliflozin 100 mg %	Canagliflozin 300 mg %
Overall population	1.5%	2.3%	3.4%
75 years of age and older <sup>†</sup>	2.6%	4.9%	8.7%
eGFR less than 60 mL/min/1.73 m <sup>2</sup> *	2.5%	4.7%	8.1%
Use of loop diuretic <sup>†</sup>	4.7%	3.2%	8.8%

\* Includes placebo and active-comparator groups

<sup>†</sup> Patients could have more than 1 of the listed risk factors

**Falls**

In a pool of nine clinical trials with mean duration of exposure to canagliflozin of 85 weeks, the proportion of patients who experienced falls was 1.3%, 1.5%, and 2.1% with comparator, canagliflozin 100 mg, and canagliflozin 300 mg, respectively. The higher risk of falls for patients treated with canagliflozin was observed within the first few weeks of treatment.

**Impairment in Renal Function**

Canagliflozin is associated with a dose-dependent increase in serum creatinine and a concomitant fall in estimated GFR (Table 3). Patients with moderate renal impairment at baseline had larger mean changes.

**Table 3: Changes in Serum Creatinine and eGFR Associated with Canagliflozin in the Pool of Four Placebo-Controlled Trials and Moderate Renal Impairment Trial**

		Placebo N=646	Canagliflozin 100 mg N=833	Canagliflozin 300 mg N=834	
Pool of Four Placebo- Controlled Trials	Baseline	Creatinine (mg/dL)	0.84	0.82	0.82
		eGFR (mL/ min/1.73 m <sup>2</sup> )	87.0	89.3	88.8
	Week 6 Change	Creatinine (mg/dL)	0.01	0.03	0.05
		eGFR (mL/min/1.73 m <sup>2</sup> )	-1.6	-3.8	-5.0
	End of Treatment Change*	Creatinine (mg/dL)	0.01	0.02	0.03
		eGFR (mL/min/1.73 m <sup>2</sup> )	-1.6	-2.3	-3.4
		Placebo N=90	Canagliflozin 100 mg N=90	Canagliflozin 300 mg N=89	
Moderate Renal Impairment Trial	Baseline	Creatinine (mg/dL)	1.81	1.62	1.63
		eGFR (mL/min/1.73 m <sup>2</sup> )	40.1	39.7	38.5
	Week 3 Change	Creatinine (mg/dL)	0.03	0.16	0.28
		eGFR (mL/min/1.73 m <sup>2</sup> )	-0.7	-4.8	-6.2
	End of Treatment Change*	Creatinine (mg/dL)	0.07	0.16	0.18
		eGFR (mL/min/1.73 m <sup>2</sup> )	-1.5	-3.6	-4.0

\* Week 26 in mITT LOCF population

**INVOKAMET® (canagliflozin and metformin hydrochloride) tablets**

In the pool of four placebo-controlled trials where patients had normal or mildly impaired baseline renal function, the proportion of patients who experienced at least one event of significant renal function decline, defined as an eGFR below 80 mL/min/1.73 m<sup>2</sup> and 30% lower than baseline, was 2.1% with placebo, 2.0% with canagliflozin 100 mg, and 4.1% with canagliflozin 300 mg. At the end of treatment, 0.5% with placebo, 0.7% with canagliflozin 100 mg, and 1.4% with canagliflozin 300 mg had a significant renal function decline.

In a trial carried out in patients with moderate renal impairment with a baseline eGFR of 30 to less than 50 mL/min/1.73 m<sup>2</sup> (mean baseline eGFR 39 mL/min/1.73 m<sup>2</sup>), the proportion of patients who experienced at least one event of significant renal function decline, defined as an eGFR 30% lower than baseline, was 6.9% with placebo, 18% with canagliflozin 100 mg, and 22.5% with canagliflozin 300 mg. At the end of treatment, 4.8% with placebo, 3.4% with canagliflozin 100 mg, and 2.2% with canagliflozin 300 mg had a significant renal function decline.

In a pooled population of patients with moderate renal impairment (N=1085) with baseline eGFR of 30 to less than 60 mL/min/1.73 m<sup>2</sup> (mean baseline eGFR 48 mL/min/1.73 m<sup>2</sup>), the overall incidence of these events was lower than in the dedicated trial but a dose-dependent increase in incident episodes of significant renal function decline compared to placebo was still observed.

Use of canagliflozin has been associated with an increased incidence of renal-related adverse reactions (e.g., increased blood creatinine, decreased glomerular filtration rate, renal impairment, and acute renal failure), particularly in patients with moderate renal impairment.

In the pooled analysis of patients with moderate renal impairment, the incidence of renal-related adverse reactions was 3.7% with placebo, 8.9% with canagliflozin 100 mg, and 9.3% with canagliflozin 300 mg. Discontinuations due to renal-related adverse events occurred in 1.0% with placebo, 1.2% with canagliflozin 100 mg, and 1.6% with canagliflozin 300 mg [see Warnings and Precautions (5.4)].

**Genital Mycotic Infections**

In the pool of four placebo-controlled clinical trials, female genital mycotic infections (e.g., vulvovaginal mycotic infection, vulvovaginal candidiasis, and vulvovaginitis) occurred in 2.8%, 10.6%, and 11.6% of females treated with placebo, canagliflozin 100 mg, and canagliflozin 300 mg, respectively. Patients with a history of genital mycotic infections were more likely to develop genital mycotic infections on canagliflozin. Female patients who developed genital mycotic infections on canagliflozin were more likely to experience recurrence and require treatment with oral or topical antifungal agents and anti-microbial agents. In females, discontinuation due to genital mycotic infections occurred in 0% and 0.7% of patients treated with placebo and canagliflozin, respectively [see Warnings and Precautions (5.8)].

In the pool of four placebo-controlled clinical trials, male genital mycotic infections (e.g., candidal balanitis, balanoposthitis) occurred in 0.7%, 4.2%, and 3.8% of males treated with placebo, canagliflozin 100 mg, and canagliflozin 300 mg, respectively. Male genital mycotic infections occurred more commonly in uncircumcised males and in males with a prior history of balanitis or balanoposthitis. Male patients who developed genital mycotic infections on canagliflozin were more likely to experience recurrent infections (22% on canagliflozin versus none on placebo), and require treatment with oral or topical antifungal agents and anti-microbial agents than patients on comparators. In males, discontinuations due to genital mycotic infections occurred in 0% and 0.5% of patients treated with placebo and canagliflozin, respectively. In the pooled analysis of 8 controlled trials, phimosis was reported in 0.3% of uncircumcised male patients treated with canagliflozin and 0.2% required circumcision to treat the phimosis [see Warnings and Precautions (5.8)].

**Hypoglycemia**

In canagliflozin clinical trials, hypoglycemia was defined as any event regardless of symptoms, where biochemical hypoglycemia was documented (any glucose value below or equal to 70 mg/dL). Severe hypoglycemia was defined as an event consistent with hypoglycemia where the patient required the assistance of another person to recover, lost consciousness, or experienced a seizure (regardless of whether biochemical documentation of a low glucose value was obtained). In individual clinical trials [see Clinical Studies (14.6)], episodes of hypoglycemia occurred at a higher rate when canagliflozin was co-administered with insulin or sulfonylureas (Table 4) [see Warnings and Precautions (5.7)].

**Table 4: Incidence of Hypoglycemia\* in Controlled Clinical Studies**

Monotherapy (26 weeks)	Placebo (N=192)	Canagliflozin 100 mg (N=195)	Canagliflozin 300 mg (N=197)
Overall [N (%)]	5 (2.6)	7 (3.6)	6 (3.0)
In Combination with Metformin (26 weeks)	Placebo + Metformin (N=183)	Canagliflozin 100 mg + Metformin (N=368)	Canagliflozin 300 mg + Metformin (N=367)
Overall [N (%)]	3 (1.6)	16 (4.3)	17 (4.6)
Severe [N (%)] <sup>†</sup>	0 (0)	1 (0.3)	1 (0.3)

**INVOKAMET® (canagliflozin and metformin hydrochloride) tablets****Table 4: Incidence of Hypoglycemia\* in Controlled Clinical Studies (continued)**

In Combination with Metformin (18 weeks) <sup>‡</sup>	Placebo (N=93)	Canagliflozin 100 mg (N=93)	Canagliflozin 300 mg (N=93)
Overall [N (%)]	3 (3.2)	4 (4.3)	3 (3.2)
In Combination with Metformin + Sulfonylurea (26 weeks)	Placebo + Sulfonylurea (N=156)	Canagliflozin 100 mg + Metformin + Sulfonylurea (N=157)	Canagliflozin 300 mg + Metformin + Sulfonylurea (N=156)
Overall [N (%)]	24 (15.4)	43 (27.4)	47 (30.1)
Severe [N (%)] <sup>†</sup>	1 (0.6)	1 (0.6)	0
In Combination with Metformin + Pioglitazone (26 weeks)	Placebo + Metformin + Pioglitazone (N=115)	Canagliflozin 100 mg + Metformin + Pioglitazone (N=113)	Canagliflozin 300 mg + Metformin + Pioglitazone (N=114)
Overall [N (%)]	3 (2.6)	3 (2.7)	6 (5.3)
In Combination with Insulin (18 weeks)	Placebo (N=565)	Canagliflozin 100 mg (N=568)	Canagliflozin 300 mg (N=567)
Overall [N (%)]	208 (36.8)	279 (49.3)	285 (49.6)
Severe [N (%)] <sup>†</sup>	14 (2.5)	10 (1.8)	16 (2.7)
In Combination with Insulin and Metformin (18 weeks) <sup>‡</sup>	Placebo (N=145)	Canagliflozin 100 mg (N=139)	Canagliflozin 300 mg (N=148)
Overall [N (%)]	66 (45.5)	58 (41.7)	70 (47.3)
Severe [N (%)] <sup>†</sup>	4 (2.8)	1 (0.7)	3 (2.0)

\* Number of patients experiencing at least one event of hypoglycemia based on either biochemically documented episodes or severe hypoglycemic events in the intent-to-treat population

<sup>†</sup> Severe episodes of hypoglycemia were defined as those where the patient required the assistance of another person to recover, lost consciousness, or experienced a seizure (regardless of whether biochemical documentation of a low glucose value was obtained)

<sup>‡</sup> Phase 2 clinical study with twice daily dosing (50 mg or 150 mg twice daily in combination with metformin)

<sup>§</sup> Subgroup of patients (N=287) from insulin substudy on canagliflozin in combination with metformin and insulin (with or without other antiglycemic agents)

**Bone Fracture**

The occurrence of bone fractures was evaluated in a pool of nine clinical trials with a mean duration of exposure to canagliflozin of 85 weeks. The incidence rates of adjudicated bone fractures were 1.1, 1.4, and 1.5 per 100 patient-years of exposure in the comparator, canagliflozin 100 mg, and canagliflozin 300 mg groups, respectively. Fractures were observed as early as 12 weeks after treatment initiation and were more likely to be low trauma (e.g., fall from no more than standing height), and affect the upper extremities [see Warnings and Precautions (5.11)].

**Metformin**

The most common adverse reactions (5% or greater incidence) due to initiation of metformin are diarrhea, nausea, vomiting, flatulence, asthenia, indigestion, abdominal discomfort, and headache.

Long-term treatment with metformin has been associated with a decrease in vitamin B<sub>12</sub>, which may very rarely result in clinically significant vitamin B<sub>12</sub> deficiency (e.g., megaloblastic anemia) [see Warnings and Precautions (5.12)].

**Laboratory and Imaging Tests****Increases in Serum Potassium**

In a pooled population of patients (N=723) with moderate renal impairment (eGFR 45 to less than 60 mL/min/1.73 m<sup>2</sup>), increases in serum potassium to greater than 5.4 mEq/L and 15% above baseline occurred in 5.3%, 5.0%, and 6.8% of patients treated with placebo, canagliflozin 100 mg, and canagliflozin 300 mg, respectively. Severe elevations (greater than or equal to 6.5 mEq/L) occurred in 0.4% of patients treated with placebo, no patients treated with canagliflozin 100 mg, and 1.3% of patients treated with canagliflozin 300 mg.

In these patients, increases in potassium were more commonly seen in those with elevated potassium at baseline. Among patients with moderate renal impairment, approximately 84% were taking medications that interfere with potassium excretion, such as potassium-sparing diuretics, angiotensin-converting-enzyme inhibitors, and angiotensin-receptor blockers [see Warnings and Precautions (5.2, 5.5) and Use in Specific Populations (8.8)].

**Increases in Serum Magnesium**

Dose-related increases in serum magnesium were observed early after initiation of canagliflozin (within 6 weeks) and remained elevated throughout treatment. In the pool of four placebo-controlled trials, the mean percent change in serum

**INVOKAMET® (canagliflozin and metformin hydrochloride) tablets**

magnesium levels was 8.1% and 9.3% with canagliflozin 100 mg and canagliflozin 300 mg, respectively, compared to -0.6% with placebo. In a trial of patients with moderate renal impairment, serum magnesium levels increased by 0.2%, 9.2%, and 14.8% with placebo, canagliflozin 100 mg, and canagliflozin 300 mg, respectively.

**Increases in Serum Phosphate**

Dose-related increases in serum phosphate levels were observed with canagliflozin. In the pool of four placebo-controlled trials, the mean percent change in serum phosphate levels were 3.6% and 5.1% with canagliflozin 100 mg and canagliflozin 300 mg, respectively, compared to 1.5% with placebo. In a trial of patients with moderate renal impairment, the mean serum phosphate levels increased by 1.2%, 5.0%, and 9.3% with placebo, canagliflozin 100 mg, and canagliflozin 300 mg, respectively.

**Increases in Low-Density Lipoprotein Cholesterol (LDL-C) and non-High-Density Lipoprotein Cholesterol (non-HDL-C)**

In the pool of four placebo-controlled trials, dose-related increases in LDL-C with canagliflozin were observed. Mean changes (percent changes) from baseline in LDL-C relative to placebo were 4.4 mg/dL (4.5%) and 8.2 mg/dL (8.0%) with canagliflozin 100 mg and canagliflozin 300 mg, respectively. The mean baseline LDL-C levels were 104 to 110 mg/dL across treatment groups [see *Warnings and Precautions* (5.12)].

Dose-related increases in non-HDL-C with canagliflozin were observed. Mean changes (percent changes) from baseline in non-HDL-C relative to placebo were 2.1 mg/dL (1.5%) and 5.1 mg/dL (3.8%) with canagliflozin 100 mg and 300 mg, respectively. The mean baseline non-HDL-C levels were 140 to 147 mg/dL across treatment groups.

**Increases in Hemoglobin**

In the pool of four placebo-controlled trials, mean changes (percent changes) from baseline in hemoglobin were -0.18 g/dL (-1.1%) with placebo, 0.47 g/dL (3.5%) with canagliflozin 100 mg, and 0.51 g/dL (3.8%) with canagliflozin 300 mg. The mean baseline hemoglobin value was approximately 14.1 g/dL across treatment groups. At the end of treatment, 0.8%, 4.0%, and 2.7% of patients treated with placebo, canagliflozin 100 mg, and canagliflozin 300 mg, respectively, had hemoglobin levels above the upper limit of normal.

**Decreases in Bone Mineral Density**

Bone mineral density (BMD) was measured by dual-energy X-ray absorptiometry in a clinical trial of 714 older adults (mean age 64 years). At 2 years, patients randomized to canagliflozin 100 mg and canagliflozin 300 mg had placebo-corrected declines in BMD at the total hip of 0.9% and 1.2%, respectively, and at the lumbar spine of 0.3% and 0.7%, respectively. Additionally, placebo-adjusted BMD declines were 0.1% at the femoral neck for both canagliflozin doses and 0.4% at the distal forearm for patients randomized to canagliflozin 300 mg. The placebo-adjusted change at the distal forearm for patients randomized to canagliflozin 100 mg was 0%.

**6.2 Postmarketing Experience**

Additional adverse reactions have been identified during postapproval use of canagliflozin. Because these reactions are reported voluntarily from a population of uncertain size, it is generally not possible to reliably estimate their frequency or establish a causal relationship to drug exposure.

**Ketoacidosis [see *Warnings and Precautions* (5.3)]**

**Acute Kidney Injury and Impairment in Renal Function [see *Warnings and Precautions* (5.4)]**

**Anaphylaxis, Angioedema [see *Warnings and Precautions* (5.9)]**

**Ursepsis and Pyelonephritis [see *Warnings and Precautions* (5.6)]**

**7 DRUG INTERACTIONS****7.1 Drug Interactions with Metformin****Carbonic Anhydrase Inhibitors**

Topiramate or other carbonic anhydrase inhibitors (e.g., zonisamide, acetazolamide or dichlorophanamide) frequently causes a decrease in serum bicarbonate and induce non-anion gap, hyperchloremic metabolic acidosis. Concomitant use of these drugs with INVOKAMET may increase the risk for lactic acidosis. Consider more frequent monitoring of these patients.

**Drugs That Reduce Metformin Clearance**

Drugs that are eliminated by renal tubular secretion (e.g. cationic drugs such as cimetidine) have the potential for interaction with metformin by competing for common renal tubular transport systems, and may increase the accumulation of metformin and the risk for lactic acidosis [see *Clinical Pharmacology* (12.3)]. Consider more frequent monitoring of these patients.

**Alcohol**

Alcohol is known to potentiate the effect of metformin on lactate metabolism. Warn patients against excessive alcohol intake while receiving INVOKAMET.

**Drugs Affecting Glycemic Control**

Certain drugs tend to produce hyperglycemia and may lead to loss of glycemic control. These drugs include the thiazides and other diuretics, corticosteroids, phenothiazines, thyroid products, estrogens, oral contraceptives, phenytoin, nicotinic acid, sympathomimetics, calcium channel blockers, and isoniazid. When such drugs are administered to a patient receiving INVOKAMET, monitor for

**INVOKAMET® (canagliflozin and metformin hydrochloride) tablets**

loss of blood glucose control. When such drugs are withdrawn from a patient receiving INVOKAMET, monitor for hypoglycemia.

**7.2 Drug Interactions with Canagliflozin****UGT Enzyme Inducers**

Rifampin: Rifampin lowered canagliflozin exposure which may reduce the efficacy of INVOKAMET. If an inducer of UGTs (e.g., rifampin, phenytoin, phenobarbital, ritonavir) must be co-administered with INVOKAMET, consider increasing the dose to canagliflozin 150 mg twice daily if patients are currently tolerating INVOKAMET with 50 mg canagliflozin twice daily, have an eGFR greater than 60 mL/min/1.73 m<sup>2</sup>, and require additional glycemic control. Consider other antihyperglycemic therapy in patients with an eGFR of 45 to less than 60 mL/min/1.73 m<sup>2</sup> receiving concurrent therapy with a UGT inducer and require additional glycemic control [see *Dosage and Administration* (2.3) and *Clinical Pharmacology* (12.3)].

**Digoxin**

Canagliflozin increased digoxin exposure. Digoxin, as a cationic drug, also has the potential to compete with metformin for common renal tubular transport systems [see *Drug Interactions* (7.1)]. Monitor patients taking INVOKAMET with concomitant digoxin for a need to adjust dose of either drug.

**Drug/Laboratory Test Interference****Positive Urine Glucose Test**

Monitoring glycemic control with urine glucose tests is not recommended in patients taking SGLT2 inhibitors as SGLT2 inhibitors increase urinary glucose excretion and will lead to positive urine glucose tests. Use alternative methods to monitor glycemic control.

**Interference with 1,5-anhydroglucitol (1,5-AG) Assay**

Monitoring glycemic control with 1,5-AG assay is not recommended as measurements of 1,5-AG are unreliable in assessing glycemic control in patients taking SGLT2 inhibitors. Use alternative methods to monitor glycemic control.

**8 USE IN SPECIFIC POPULATIONS****8.1 Pregnancy****Risk Summary**

Based on animal data showing adverse renal effects, INVOKAMET is not recommended during the second and third trimesters of pregnancy.

Limited data with INVOKAMET or canagliflozin in pregnant women are not sufficient to determine a drug-associated risk for major birth defects or miscarriage. Published studies with metformin use during pregnancy have not reported a clear association with metformin and major birth defect or miscarriage risk [see *Data*]. There are risks to the mother and fetus associated with poorly controlled diabetes in pregnancy [see *Clinical Considerations*].

In animal studies, adverse renal pelvic and tubule dilatations that were not reversible were observed in rats when canagliflozin was administered at an exposure 0.5-times the 300 mg clinical dose, based on AUC during a period of renal development corresponding to the late second and third trimesters of human pregnancy. No adverse developmental effects were observed when metformin was administered to pregnant Sprague Dawley rats and rabbits during the period of organogenesis at doses up to 2- and 6-times, respectively, a 2000 mg clinical dose, based on body surface area [see *Data*].

The estimated background risk of major birth defects is 6-10% in women with pre-gestational diabetes with an HbA1c >7 and has been reported to be as high as 20-25% in women with a HbA1c >10. The estimated background risk of miscarriage for the indicated population is unknown. In the U.S. general population, the estimated background risk of major birth defects and miscarriage in clinically recognized pregnancies is 2-4% and 15-20%, respectively.

**Clinical Considerations****Disease-associated maternal and/or embryo/fetal risk**

Poorly controlled diabetes in pregnancy increases the maternal risk for diabetic ketoacidosis, pre-eclampsia, spontaneous abortions, preterm delivery, stillbirth and delivery complications. Poorly controlled diabetes increases the fetal risk for major birth defects, stillbirth, and macrosomia related morbidity.

**Data****Human Data**

Published data from post-marketing studies have not reported a clear association with metformin and major birth defects, miscarriage, or adverse maternal or fetal outcomes when metformin was used during pregnancy. However, these studies cannot definitely establish the absence of any metformin-associated risk because of methodological limitations, including small sample size and inconsistent comparator groups.

**Animal Data****Canagliflozin**

Canagliflozin dosed directly to juvenile rats from postnatal day (PND) 21 until PND 90 at doses of 4, 20, 65, or 100 mg/kg increased kidney weights and dose dependently increased the incidence and severity of renal pelvic and tubular dilatation at all doses tested. Exposure at the lowest dose was greater than or equal to 0.5-times the 300 mg clinical dose, based on AUC. These outcomes occurred with drug exposure during periods of renal development in rats that correspond to the late second and third trimester of human renal development.

**INVOKAMET® (canagliflozin and metformin hydrochloride) tablets**

The renal pelvic dilatations observed in juvenile animals did not fully reverse within a 1 month recovery period.

In embryo-fetal development studies in rats and rabbits, canagliflozin was administered for intervals coinciding with the first trimester period of organogenesis in humans. No developmental toxicities independent of maternal toxicity were observed when canagliflozin was administered at doses up to 100 mg/kg in pregnant rats and 180 mg/kg in pregnant rabbits during embryonic organogenesis or during a study in which maternal rats were dosed from gestation day (GD) 6 through PND 21, yielding exposures up to approximately 18-times the 300 mg clinical dose, based on AUC.

**Metformin Hydrochloride**

Metformin hydrochloride did not cause adverse developmental effects when administered to pregnant Sprague Dawley rats and rabbits up to 600 mg/kg/day during the period of organogenesis. This represents an exposure of about 2- and 6-times a 2000 mg clinical dose based on body surface area (mg/m<sup>2</sup>) for rats and rabbits, respectively.

**Canagliflozin and Metformin**

No adverse developmental effects were observed when canagliflozin and metformin were co-administered to pregnant rats during the period of organogenesis at exposures up to 11 and 13 times, respectively, the 300 mg and 2000 mg clinical doses of canagliflozin and metformin based on AUC.

**8.2 Lactation****Risk Summary**

There is no information regarding the presence of INVOKAMET or canagliflozin in human milk, the effects on the breastfed infant, or the effects on milk production. Limited published studies report that metformin is present in human milk [see Data]. However, there is insufficient information on the effects of metformin on the breastfed infant and no available information on the effects of metformin on milk production. Canagliflozin is present in the milk of lactating rats [see Data]. Since human kidney maturation occurs *in utero* and during the first 2 years of life when lactational exposure may occur, there may be risk to the developing human kidney.

Because of the potential for serious adverse reactions in a breastfed infant, advise women that use of INVOKAMET is not recommended while breastfeeding.

**Data****Human Data**

Published clinical lactation studies report that metformin is present in human milk which resulted in infant doses approximately 0.11% to 1% of the maternal weight-adjusted dosage and a milk/plasma ratio ranging between 0.13 and 1. However, the studies were not designed to definitely establish the risk of use of metformin during lactation because of small sample size and limited adverse event data collected in infants.

**Animal Data**

Radio-labeled canagliflozin administered to lactating rats on day 13 post-partum was present at a milk/plasma ratio of 1.40, indicating that canagliflozin and its metabolites are transferred into milk at a concentration comparable to that in plasma. Juvenile rats directly exposed to canagliflozin showed a risk to the developing kidney (renal pelvic and tubular dilatations) during maturation.

**8.3 Females and Males of Reproductive Potential**

Discuss the potential for unintended pregnancy with premenopausal women as therapy with metformin may result in ovulation in some anovulatory women.

**8.4 Pediatric Use**

Safety and effectiveness of INVOKAMET in pediatric patients under 10 years of age have not been established.

**8.5 Geriatric Use****INVOKAMET**

Because renal function abnormalities can occur after initiating canagliflozin, metformin is substantially excreted by the kidney, and aging can be associated with reduced renal function, monitor renal function more frequently after initiating INVOKAMET in the elderly and then adjust dose based on renal function [see Dosage and Administration (2.2) and Warnings and Precautions (5.1, 5.3)].

**Canagliflozin**

Two thousand thirty-four (2034) patients 65 years and older, and 345 patients 75 years and older were exposed to canagliflozin in nine clinical studies of canagliflozin. Of these patients, 1334 patients 65 years and older and 181 patients 75 years and older were exposed to the combination of canagliflozin and metformin [see Clinical Studies (14)]. Patients 65 years and older had a higher incidence of adverse reactions related to reduced intravascular volume with canagliflozin (such as hypotension, postural dizziness, orthostatic hypotension, syncope, and dehydration), particularly with the 300 mg daily dose, compared to younger patients; a more prominent increase in the incidence was seen in patients who were 75 years and older [see Dosage and Administration (2.1) and Adverse Reactions (6.1)]. Smaller reductions in HbA1C with canagliflozin relative to placebo were seen in older (65 years and older; -0.61% with canagliflozin 100 mg and -0.74% with canagliflozin 300 mg relative to placebo) compared to younger patients (-0.72% with canagliflozin 100 mg and -0.87% with canagliflozin 300 mg relative to placebo).

**INVOKAMET® (canagliflozin and metformin hydrochloride) tablets****Metformin**

Controlled clinical studies of metformin did not include sufficient numbers of elderly patients to determine whether they respond differently from younger patients, although other reported clinical experience has not identified differences in responses between the elderly and younger patients. The initial and maintenance dosing of metformin should be conservative in patients with advanced age due to the potential for decreased renal function in this population. Any dose adjustment should be based on a careful assessment of renal function [see Contraindications (4), Warnings and Precautions (5.3), and Clinical Pharmacology (12.3)].

**8.6 Renal Impairment****Canagliflozin**

The efficacy and safety of canagliflozin were evaluated in a study that included patients with moderate renal impairment (eGFR 30 to less than 50 mL/min/1.73 m<sup>2</sup>). These patients had less overall glycemic efficacy and had a higher occurrence of adverse reactions related to reduced intravascular volume, renal-related adverse reactions, and decreases in eGFR compared to patients with mild renal impairment or normal renal function (eGFR greater than or equal to 60 mL/min/1.73 m<sup>2</sup>). Dose-related, transient mean increases in serum potassium were observed early after initiation of canagliflozin (i.e., within 3 weeks) in this trial. Increases in serum potassium of greater than 5.4 mEq/L and 15% above baseline occurred in 16.1%, 12.4%, and 27.0% of patients treated with placebo, canagliflozin 100 mg, and canagliflozin 300 mg, respectively. Severe elevations (greater than or equal to 6.5 mEq/L) occurred in 1.1%, 2.2%, and 2.2% of patients treated with placebo, canagliflozin 100 mg, and canagliflozin 300 mg, respectively [see Dosage and Administration (2.2), Contraindications (4), Warnings and Precautions (5.1, 5.3, 5.4), and Adverse Reactions (6.1)].

The efficacy and safety of canagliflozin have not been established in patients with severe renal impairment (eGFR less than 30 mL/min/1.73 m<sup>2</sup>), with ESRD, or receiving dialysis. Canagliflozin is not expected to be effective in these patient populations [see Contraindications (4) and Clinical Pharmacology (12.3)].

**8.7 Hepatic Impairment**

Use of metformin in patients with hepatic impairment has been associated with some cases of lactic acidosis. INVOKAMET is not recommended in patients with hepatic impairment. [see Warnings and Precautions (5.1)]

**10 OVERDOSAGE**

In the event of an overdose with INVOKAMET, contact the Poison Control Center. Employ the usual supportive measures (e.g., remove unabsorbed material from the gastrointestinal tract, employ clinical monitoring, and institute supportive treatment) as dictated by the patient's clinical status. Canagliflozin was negligibly removed during a 4-hour hemodialysis session. Canagliflozin is not expected to be dialyzable by peritoneal dialysis. Metformin is dialyzable with a clearance of up to 170 mL/min under good hemodynamic conditions. Therefore, hemodialysis may be useful partly for removal of accumulated metformin from patients in whom INVOKAMET overdosage is suspected.

**Canagliflozin**

There were no reports of overdose during the clinical development program of canagliflozin.

**Metformin**


Overdose of metformin hydrochloride has occurred, including ingestion of amounts greater than 50 grams. Hypoglycemia was reported in approximately 10% of cases, but no causal association with metformin hydrochloride has been established. Lactic acidosis has been reported in approximately 32% of metformin overdose cases [see Warnings and Precautions (5.1)].

**11 DESCRIPTION**

INVOKAMET (canagliflozin and metformin hydrochloride) tablets contain two oral antihyperglycemic drugs used in the management of type 2 diabetes: canagliflozin and metformin hydrochloride.

**Canagliflozin**


Canagliflozin is an inhibitor of sodium-glucose co-transporter 2 (SGLT2), the transporter responsible for reabsorbing the majority of glucose filtered by the kidney. Canagliflozin is chemically known as (1S)-1,5-anhydro-1-[3-[[5-(4-fluorophenyl)-2-thienyl]methyl]-4-methylphenyl]-D-glucitol hemihydrate and its molecular formula and weight are C<sub>24</sub>H<sub>28</sub>F<sub>2</sub>O<sub>5</sub>•1/2 H<sub>2</sub>O and 453.53, respectively. The structural formula for canagliflozin is:



Canagliflozin is practically insoluble in aqueous media from pH 1.1 to 12.9.

**INVOKAMET® (canagliflozin and metformin hydrochloride) tablets****Metformin Hydrochloride**

Metformin hydrochloride is not chemically or pharmacologically related to any other classes of oral antihyperglycemic agents. Metformin hydrochloride is chemically known as 1,1-Dimethylbiguanide hydrochloride and its molecular formula and weight are  $C_4H_{11}N_5 \cdot HCl$  and 165.62, respectively. The structural formula for metformin hydrochloride is:

**INVOKAMET**

INVOKAMET is supplied as film-coated tablets for oral administration. Each 50 mg/500 mg tablet and 50 mg/1,000 mg tablet contains 51 mg of canagliflozin equivalent to 50 mg canagliflozin (anhydrous) and 500 mg or 1,000 mg metformin hydrochloride. Each 150 mg/500 mg tablet and 150 mg/1,000 mg tablet contains 153 mg of canagliflozin equivalent to 150 mg canagliflozin (anhydrous) and 500 mg or 1,000 mg metformin hydrochloride.

Inactive ingredients of the core tablet are croscarmellose sodium, hypromellose, magnesium stearate, and microcrystalline cellulose. The magnesium stearate is vegetable-sourced. The tablets are finished with a commercially available film-coating consisting of the following excipients: Macrolog/PEG, polyvinyl alcohol (partially hydrolyzed), talc, titanium dioxide, iron oxide yellow, (50 mg/1,000 mg and 150 mg/500 mg tablets only), iron oxide red, (50 mg/1,000 mg, 150 mg/500 mg and 150 mg/1,000 mg tablets only), and iron oxide black (150 mg/1,000 mg tablets only).

**12 CLINICAL PHARMACOLOGY****12.1 Mechanism of Action****INVOKAMET**

INVOKAMET (canagliflozin and metformin hydrochloride) combines two oral antihyperglycemic agents with complementary mechanisms of action to improve glycemic control in patients with type 2 diabetes: canagliflozin, a sodium-glucose co-transporter 2 (SGLT2) inhibitor, and metformin hydrochloride, a member of the biguanide class.

**Canagliflozin**

Sodium-glucose co-transporter 2 (SGLT2), expressed in the proximal renal tubules, is responsible for the majority of the reabsorption of filtered glucose from the tubular lumen. Canagliflozin is an inhibitor of SGLT2. By inhibiting SGLT2, canagliflozin reduces reabsorption of filtered glucose and lowers the renal threshold for glucose ( $RT_G$ ), and thereby increases urinary glucose excretion (UGE).

**Metformin**

Metformin is an antihyperglycemic agent which improves glucose tolerance in patients with type 2 diabetes, lowering both basal and postprandial plasma glucose. Metformin decreases hepatic glucose production, decreases intestinal absorption of glucose, and improves insulin sensitivity by increasing peripheral glucose uptake and utilization. Metformin does not produce hypoglycemia in either patients with type 2 diabetes or normal patients except in special circumstances [see *Warnings and Precautions (5.7)*] and does not cause hyperinsulinemia. With metformin therapy, insulin secretion remains unchanged while fasting insulin levels and day-long plasma insulin response may actually decrease.

**12.2 Pharmacodynamics****Canagliflozin**

Following single and multiple oral doses of canagliflozin in patients with type 2 diabetes, dose-dependent decreases in  $RT_G$  and increases in urinary glucose excretion were observed. From a starting  $RT_G$  value of approximately 240 mg/dL, canagliflozin at 100 mg and 300 mg once daily suppressed  $RT_G$  throughout the 24-hour period. Maximal suppression of mean  $RT_G$  over the 24-hour period was seen with the 300 mg daily dose to approximately 70 to 90 mg/dL in patients with type 2 diabetes in Phase 1 studies. The reductions in  $RT_G$  led to increases in mean UGE of approximately 100 g/day in patients with type 2 diabetes treated with either 100 mg or 300 mg of canagliflozin. The 24-h mean  $RT_G$  at steady state was similar following once daily and twice daily dosing regimens at the same total daily dose of 100 mg or 300 mg. In patients with type 2 diabetes given 100 to 300 mg once daily over a 18-day dosing period, reductions in  $RT_G$  and increases in urinary glucose excretion were observed over the dosing period. In this study, plasma glucose declined in a dose-dependent fashion within the first day of dosing.

**Cardiac Electrophysiology**

In a randomized, double-blind, placebo-controlled, active-comparator, 4-way crossover study, 60 healthy subjects were administered a single oral dose of canagliflozin 300 mg, canagliflozin 1,200 mg (4 times the maximum recommended dose), moxifloxacin, and placebo. No meaningful changes in QTc interval were observed with either the recommended dose of 300 mg or the 1,200 mg dose.

**INVOKAMET® (canagliflozin and metformin hydrochloride) tablets****12.3 Pharmacokinetics****INVOKAMET**

The results of a bioequivalence study in healthy subjects demonstrated that INVOKAMET 50 mg/500 mg, 50 mg/1,000 mg, 150 mg/500 mg and 150 mg/1,000 mg combination tablets are bioequivalent to co-administration of corresponding doses of canagliflozin and metformin hydrochloride as individual tablets under fed conditions.

Administration of INVOKAMET 150 mg/1,000 mg fixed-dose combination with food resulted in no change in overall exposure of canagliflozin. There was no change in metformin AUC; however, the mean peak plasma concentration of metformin was decreased by 16% when administered with food. A delayed time to peak plasma concentration was observed for both components (a delay of 2 hours for canagliflozin and 1 hour for metformin) under fed conditions. These changes are not likely to be clinically meaningful.

**Canagliflozin**

The pharmacokinetics of canagliflozin is essentially similar in healthy subjects and patients with type 2 diabetes. Following single-dose oral administration of 100 mg and 300 mg of canagliflozin, peak plasma concentrations (median  $T_{max}$ ) of canagliflozin occurs within 1 to 2 hours post-dose. Plasma  $C_{max}$  and AUC of canagliflozin increased in a dose-proportional manner from 50 mg to 300 mg. The apparent terminal half-life ( $t_{1/2}$ ) was 10.6 hours and 13.1 hours for the 100 mg and 300 mg doses, respectively. Steady-state was reached after 4 to 5 days of once-daily dosing with canagliflozin 100 mg to 300 mg. Canagliflozin does not exhibit time-dependent pharmacokinetics and accumulated in plasma up to 36% following multiple doses of 100 mg and 300 mg. The mean systemic exposure (AUC) at steady state was similar following once daily and twice daily dosing regimens at the same total daily dose of 100 mg or 300 mg.

**Absorption****Canagliflozin**

The mean absolute oral bioavailability of canagliflozin is approximately 65%.

**Metformin**

The absolute bioavailability of a metformin hydrochloride 500 mg tablet given under fasting conditions is approximately 50% to 80%. Studies using single oral doses of metformin hydrochloride 500 to 1,500 mg, and 850 to 2,550 mg, indicate that there is a lack of dose proportionality with increasing doses, which is due to decreased absorption rather than an alteration in elimination.

**Distribution****Canagliflozin**

The mean steady-state volume of distribution of canagliflozin following a single intravenous infusion in healthy subjects was 83.5 L, suggesting extensive tissue distribution. Canagliflozin is extensively bound to proteins in plasma (99%), mainly to albumin. Protein binding is independent of canagliflozin plasma concentrations. Plasma protein binding is not meaningfully altered in patients with renal or hepatic impairment.

**Metformin**

The apparent volume of distribution (V/F) of metformin following single oral doses of metformin hydrochloride 850 mg tablets averaged  $654 \pm 358$  L. Metformin is negligibly bound to plasma proteins, in contrast to sulfonylureas, which are more than 90% protein bound. Metformin partitions into erythrocytes, most likely as a function of time. At usual clinical doses and dosing schedules of metformin tablets, steady-state plasma concentrations of metformin are reached within 24 to 48 hours and are generally less than 1 mcg/mL. During controlled clinical trials of metformin, maximum metformin plasma levels did not exceed 5 mcg/mL, even at maximum doses.

**Metabolism****Canagliflozin**

O-glucuronidation is the major metabolic elimination pathway for canagliflozin, which is mainly glucuronidated by UGT1A9 and UGT2B4 to two inactive O-glucuronide metabolites. CYP3A4-mediated (oxidative) metabolism of canagliflozin is minimal (approximately 7%) in humans.

**Metformin**

Intravenous single-dose studies in normal subjects demonstrate that metformin is excreted unchanged in the urine and does not undergo hepatic metabolism (no metabolites have been identified in humans) or biliary excretion.

**Excretion****Canagliflozin**

Following administration of a single oral [ $^{14}C$ ] canagliflozin dose to healthy subjects, 41.5%, 7.0%, and 3.2% of the administered radioactive dose was recovered in feces as canagliflozin, a hydroxylated metabolite, and an O-glucuronide metabolite, respectively. Enterohepatic circulation of canagliflozin was negligible.

Approximately 33% of the administered radioactive dose was excreted in urine, mainly as O-glucuronide metabolites (30.5%). Less than 1% of the dose was excreted as unchanged canagliflozin in urine. Renal clearance of canagliflozin 100 mg and 300 mg doses ranged from 1.30 to 1.55 mL/min.

Mean systemic clearance of canagliflozin was approximately 192 mL/min in healthy subjects following intravenous administration.

**INVOKAMET® (canagliflozin and metformin hydrochloride) tablets****Metformin**

Renal clearance is approximately 3.5 times greater than creatinine clearance, which indicates that tubular secretion is the major route of metformin elimination. Following oral administration, approximately 90% of the absorbed drug is eliminated via the renal route within the first 24 hours, with a plasma elimination half-life of approximately 6.2 hours. In blood, the elimination half-life is approximately 17.8 hours, suggesting that the erythrocyte mass may be a compartment of distribution.

**Specific Populations**

Studies characterizing the pharmacokinetics of canagliflozin and metformin after administration of INVOKAMET were not conducted in patients with renal and hepatic impairment. Descriptions of the individual components in this patient population are described below.

**Renal Impairment****Canagliflozin**

A single-dose, open-label study evaluated the pharmacokinetics of canagliflozin 200 mg in subjects with varying degrees of renal impairment (classified using the MDRD-eGFR formula) compared to healthy subjects.

Renal impairment did not affect the  $C_{max}$  of canagliflozin. Compared to healthy subjects (N=3; eGFR greater than or equal to 90 mL/min/1.73 m<sup>2</sup>), plasma AUC of canagliflozin was increased by approximately 15%, 29%, and 53% in subjects with mild (N=10), moderate (N=8), and severe (N=10) renal impairment, respectively, (eGFR 60 to less than 90, 30 to less than 60, and 15 to less than 30 mL/min/1.73 m<sup>2</sup>, respectively) but was similar for ESRD (N=8) subjects and healthy subjects. Increases in canagliflozin AUC of this magnitude are not considered clinically relevant. The pharmacodynamic response to canagliflozin declines with increasing severity of renal impairment [see **Contraindications (4)** and **Warnings and Precautions (5.4)**].

Canagliflozin was negligibly removed by hemodialysis.

**Metformin**

In patients with decreased renal function (based on measured creatinine clearance), the plasma and blood half-life of metformin is prolonged and the renal clearance is decreased in proportion to the decrease in creatinine clearance [see **Contraindications (4)** and **Warnings and Precautions (5.4)**].

**Hepatic Impairment****Canagliflozin**

Relative to subjects with normal hepatic function, the geometric mean ratios for  $C_{max}$  and AUC<sub>0-24</sub> of canagliflozin were 107% and 110%, respectively, in subjects with Child-Pugh class A (mild hepatic impairment) and 96% and 111%, respectively, in subjects with Child-Pugh class B (moderate hepatic impairment) following administration of a single 300 mg dose of canagliflozin.

These differences are not considered to be clinically meaningful. There is no clinical experience in patients with Child-Pugh class C (severe) hepatic impairment [see **Warnings and Precautions (5.1)**].

**Metformin**

No pharmacokinetic studies of metformin have been conducted in patients with hepatic insufficiency [see **Warnings and Precautions (5.1)**].

**Pharmacokinetic Effects of Age, Body Mass Index (BMI)/Weight, Gender and Race****Canagliflozin**

Based on the population PK analysis with data collected from 1526 subjects, age, body mass index (BMI)/weight, gender, and race do not have a clinically meaningful effect on the pharmacokinetics of canagliflozin [see **Use in Specific Populations (8.5)**].

**Metformin**

Metformin pharmacokinetic parameters did not differ significantly between normal subjects and patients with type 2 diabetes when analyzed according to gender.

No studies of metformin pharmacokinetic parameters according to race have been performed.

**Geriatric****INVOKAMET**

Studies characterizing the pharmacokinetics of canagliflozin and metformin after administration of INVOKAMET in geriatric patients have not been performed [see **Warnings and Precautions (5.1, 5.4)** and **Use in Specific Populations (8.5)**].

**Canagliflozin**

Age had no clinically meaningful effect on the pharmacokinetics of canagliflozin based on a population pharmacokinetic analysis [see **Adverse Reactions (6.1)** and **Use in Specific Populations (8.5)**].

**Metformin**

Limited data from controlled pharmacokinetic studies of metformin in healthy elderly subjects suggest that total plasma clearance of metformin is decreased, the half-life is prolonged, and  $C_{max}$  is increased, compared with healthy young subjects. From these data, it appears that the change in metformin pharmacokinetics with aging is primarily accounted for by a change in renal function.

**Pediatric**

Studies characterizing the pharmacokinetics of canagliflozin and metformin after administration of INVOKAMET in pediatric patients have not been performed.

**INVOKAMET® (canagliflozin and metformin hydrochloride) tablets****Drug-Drug Interactions****INVOKAMET**

Pharmacokinetic drug interaction studies with INVOKAMET have not been performed; however, such studies have been conducted with the individual components canagliflozin and metformin hydrochloride.

Co-administration of multiple doses of canagliflozin (300 mg) and metformin (2,000 mg) given once daily did not meaningfully alter the pharmacokinetics of either canagliflozin or metformin in healthy subjects.

**Canagliflozin****In Vitro Assessment of Drug Interactions**

Canagliflozin did not induce CYP450 enzyme expression (3A4, 2C9, 2C19, 2B6, and 1A2) in cultured human hepatocytes. Canagliflozin did not inhibit the CYP450 isoenzymes (1A2, 2A6, 2C19, 2D6, or 2E1) and weakly inhibited CYP2B6, CYP2C8, CYP2C9, and CYP3A4 based on *in vitro* studies with human hepatic microsomes. Canagliflozin is a weak inhibitor of P-gp.

Canagliflozin is also a substrate of drug transporters P-glycoprotein (P-gp) and MRP2.

**In Vivo Assessment of Drug Interactions****Table 5: Effect of Co-Administered Drugs on Systemic Exposures of Canagliflozin**

Co-Administered Drug	Dose of Co-Administered Drug <sup>†</sup>	Dose of Canagliflozin <sup>*</sup>	Geometric Mean Ratio (Ratio With/Without Co-Administered Drug) No Effect = 1.0	
			AUC <sup>†</sup> (90% CI)	$C_{max}$ (90% CI)
<b>See Drug Interactions (7.2) for the clinical relevance of the following:</b>				
Rifampin	600 mg QD for 8 days	300 mg	0.49 (0.44; 0.54)	0.72 (0.61; 0.84)
<b>No dose adjustments of canagliflozin required for the following:</b>				
Cyclosporine	400 mg	300 mg QD for 8 days	1.23 (1.19; 1.27)	1.01 (0.91; 1.11)
Ethinyl estradiol and levonorgestrel	0.03 mg ethinyl estradiol and 0.15 mg levonorgestrel	200 mg QD for 6 days	0.91 (0.88; 0.94)	0.92 (0.84; 0.99)
Hydrochlorothiazide	25 mg QD for 35 days	300 mg QD for 7 days	1.12 (1.08; 1.17)	1.15 (1.08; 1.25)
Metformin	2,000 mg	300 mg QD for 8 days	1.10 (1.05; 1.15)	1.05 (0.96; 1.16)
Probenecid	500 mg BID for 3 days	300 mg QD for 17 days	1.21 (1.16; 1.25)	1.13 (1.00; 1.28)

<sup>\*</sup> Single dose unless otherwise noted

<sup>†</sup> AUC<sub>0-24</sub> for drugs given as a single dose and AUC<sub>24h</sub> for drugs given as multiple doses

QD = once daily; BID = twice daily

**Table 6: Effect of Canagliflozin on Systemic Exposure of Co-Administered Drugs**

Co-Administered Drug	Dose of Co-Administered Drug <sup>*</sup>	Dose of Canagliflozin <sup>*</sup>	Geometric Mean Ratio (Ratio With/Without Co-Administered Drug) No Effect = 1.0		
			AUC <sup>†</sup> (90% CI)	$C_{max}$ (90% CI)	
<b>See Drug Interactions (7.2) for the clinical relevance of the following:</b>					
Digoxin	0.5 mg QD first day followed by 0.25 mg QD for 6 days	300 mg QD for 7 days	1.20 (1.12; 1.28)	1.36 (1.21; 1.53)	
<b>No dose adjustments of co-administered drug required for the following:</b>					
Acetaminophen	1,000 mg	300 mg BID for 25 days	1.06 <sup>‡</sup> (0.98; 1.14)	1.00 (0.82; 1.09)	
Ethinyl estradiol and levonorgestrel	0.03 mg ethinyl estradiol and 0.15 mg levonorgestrel	200 mg QD for 6 days	ethinyl estradiol	1.07 (0.99; 1.15)	1.22 (1.10; 1.35)
			levonorgestrel	1.06 (1.00; 1.13)	1.22 (1.11; 1.35)
Glyburide	1.25 mg	200 mg QD for 6 days	glyburide	1.02 (0.98; 1.07)	0.93 (0.85; 1.01)
			3-cis-hydroxy-glyburide	1.01 (0.98; 1.07)	0.99 (0.91; 1.08)
			4-trans-hydroxy-glyburide	1.03 (0.97; 1.09)	0.96 (0.88; 1.04)
Hydrochlorothiazide	25 mg QD for 35 days	300 mg QD for 7 days	0.99 (0.95; 1.04)	0.94 (0.87; 1.01)	
Metformin	2,000 mg	300 mg QD for 8 days	1.20 (1.08; 1.34)	1.06 (0.93; 1.20)	

**INVOKAMET® (canagliflozin and metformin hydrochloride) tablets****Table 6: Effect of Canagliflozin on Systemic Exposure of Co-Administered Drugs (continued)**

Co-Administered Drug	Dose of Co-Administered Drug*	Dose of Canagliflozin*	Geometric Mean Ratio (Ratio With/Without Co-Administered Drug) No Effect = 1.0		
			AUC <sup>†</sup> (90% CI)	C <sub>max</sub> (90% CI)	
Simvastatin	40 mg	300 mg QD for 7 days	simvastatin	1.12 (0.94; 1.33)	1.09 (0.91; 1.31)
			simvastatin acid	1.18 (1.03; 1.35)	1.26 (1.10; 1.45)
Warfarin	30 mg	300 mg QD for 12 days	(R)-warfarin	1.01 (0.96; 1.06)	1.03 (0.94; 1.13)
			(S)-warfarin	1.08 (1.00; 1.12)	1.01 (0.90; 1.13)
			INR	1.00 (0.98; 1.03)	1.05 (0.99; 1.12)

\* Single dose unless otherwise noted

† AUC<sub>0-∞</sub> for drugs given as a single dose and AUC<sub>0-24h</sub> for drugs given as multiple doses‡ AUC<sub>0-12h</sub>

QD = once daily; BID = twice daily; INR = International Normalized Ratio

**Metformin****Table 7: Effect of Co-Administered Drugs on Plasma Metformin Systemic Exposure**

Co-Administered Drug	Dose of Co-Administered Drug*	Dose of Metformin*	Geometric Mean Ratio (Ratio With/Without Co-Administered Drug) No Effect = 1.00	
			AUC <sup>†</sup>	C <sub>max</sub>
<b>No dose adjustments required for the following:</b>				
Glyburide	5 mg	500 mg <sup>‡</sup>	0.98 <sup>§</sup>	0.99 <sup>§</sup>
Furosemide	40 mg	850 mg	1.09 <sup>§</sup>	1.22 <sup>§</sup>
Nifedipine	10 mg	850 mg	1.18	1.21
Propranolol	40 mg	850 mg	0.90	0.94
Ibuprofen	400 mg	850 mg	1.05 <sup>§</sup>	1.07 <sup>§</sup>
<b>Drugs that are eliminated by renal tubular secretion increase the accumulation of metformin [see Warnings and Precautions (5) and Drug Interactions (7)]</b>				
Cimetidine	400 mg	850 mg	1.40	1.61
<b>Carbonic anhydrase inhibitors may cause metabolic acidosis [see Warnings and Precautions (5) and Drug Interactions (7)]</b>				
Topiramate <sup>¶</sup>	100 mg	500 mg	1.25 <sup>§</sup>	1.18

\* Single dose unless otherwise noted

† AUC = AUC<sub>0-∞</sub>

‡ Metformin hydrochloride extended-release tablets 500 mg

§ Ratio of arithmetic means

¶ Healthy volunteer study at steady state with topiramate 100 mg every 12 hours and metformin 500 mg every 12 hours for 7 days. Study conducted to assess pharmacokinetics only

†† Steady state AUC<sub>0-12h</sub>**Table 8: Effect of Metformin on Co-Administered Drug Systemic Exposures**

Co-Administered Drug	Dose of Co-Administered Drug*	Dose of Metformin*	Geometric Mean Ratio (Ratio With/Without Co-Administered Drug) No Effect = 1.00	
			AUC <sup>†</sup>	C <sub>max</sub>
<b>No dose adjustments required for the following:</b>				
Glyburide	5 mg	500 mg <sup>‡</sup>	0.78 <sup>§</sup>	0.63 <sup>§</sup>
Furosemide	40 mg	850 mg	0.87 <sup>§</sup>	0.69 <sup>§</sup>
Nifedipine	10 mg	850 mg	1.10 <sup>§</sup>	1.08
Propranolol	40 mg	850 mg	1.01 <sup>§</sup>	0.94
Ibuprofen	400 mg	850 mg	0.97 <sup>§</sup>	1.01 <sup>§</sup>
Cimetidine	400 mg	850 mg	0.95 <sup>§</sup>	1.01

\* Single dose unless otherwise noted

† AUC = AUC<sub>0-∞</sub>‡ AUC<sub>0-24h</sub> reported

§ Ratio of arithmetic means, p-value of difference &lt;0.05

†† Ratio of arithmetic means.

**INVOKAMET® (canagliflozin and metformin hydrochloride) tablets**

Metformin is negligibly bound to plasma proteins and is, therefore, less likely to interact with highly protein-bound drugs such as salicylates, sulfonamides, chloramphenicol, and probenecid.

**13 NONCLINICAL TOXICOLOGY****13.1 Carcinogenesis, Mutagenesis, Impairment of Fertility****INVOKAMET**

No animal studies have been conducted with the combined products in INVOKAMET to evaluate carcinogenesis, mutagenesis, or impairment of fertility. The following data are based on findings in studies with canagliflozin and metformin individually.

**Canagliflozin****Carcinogenesis**

Carcinogenicity was evaluated in 2-year studies conducted in CD1 mice and Sprague-Dawley rats. Canagliflozin did not increase the incidence of tumors in mice dosed at 10, 30, or 100 mg/kg (less than or equal to 14 times exposure from a 300 mg clinical dose).

Testicular Leydig cell tumors, considered secondary to increased luteinizing hormone (LH), increased significantly in male rats at all doses tested (10, 30, and 100 mg/kg). In a 12-week clinical study, LH did not increase in males treated with canagliflozin.

Renal tubular adenoma and carcinoma increased significantly in male and female rats dosed at 100 mg/kg, or approximately 12-times exposure from a 300 mg clinical dose. Also, adrenal pheochromocytoma increased significantly in males and numerically in females dosed at 100 mg/kg. Carbohydrate malabsorption associated with high doses of canagliflozin was considered a necessary proximal event in the emergence of renal and adrenal tumors in rats. Clinical studies have not demonstrated carbohydrate malabsorption in humans at canagliflozin doses of up to 2-times the recommended clinical dose of 300 mg.

**Mutagenesis**

Canagliflozin was not mutagenic with or without metabolic activation in the Ames assay. Canagliflozin was mutagenic in the *in vitro* mouse lymphoma assay with but not without metabolic activation. Canagliflozin was not mutagenic or clastogenic in an *in vivo* oral micronucleus assay in rats and an *in vivo* oral Comet assay in rats.

**Metformin****Carcinogenesis**

Long-term carcinogenicity studies have been performed in rats (dosing duration of 104 weeks) and mice (dosing duration of 91 weeks) at doses up to and including 900 mg/kg/day and 1500 mg/kg/day, respectively. These doses are both approximately 4 times the maximum recommended human daily dose of 2000 mg based on body surface area comparisons. No evidence of carcinogenicity with metformin was found in either male or female mice. Similarly, there was no tumorigenic potential observed with metformin in male rats. There was, however, an increased incidence of benign stromal uterine polyps in female rats treated with 900 mg/kg/day.

**Mutagenesis**

There was no evidence of a mutagenic potential of metformin in the following *in vitro* tests: Ames test (*S. typhimurium*), gene mutation test (mouse lymphoma cells), or chromosomal aberrations test (human lymphocytes). Results in the *in vivo* mouse micronucleus test were also negative.

**Impairment of Fertility**

Canagliflozin had no effects on the ability of rats to mate and sire or maintain a litter up to the high dose of 100 mg/kg (approximately 14 times and 18 times the 300 mg clinical dose in males and females, respectively), although there were minor alterations in a number of reproductive parameters (decreased sperm velocity, increased number of abnormal sperm, slightly fewer corpora lutea, fewer implantation sites, and smaller litter sizes) at the highest dosage administered.

Fertility of male or female rats was unaffected by metformin when administered at doses as high as 600 mg/kg/day, which is approximately 3 times the maximum recommended human daily dose based on body surface area comparisons.

**14 CLINICAL STUDIES**

Canagliflozin has been studied in combination with metformin alone, metformin and sulfonylurea, metformin and a thiazolidinedione (i.e. pioglitazone), and metformin and insulin (with or without other anti-hyperglycemic agents). The efficacy of canagliflozin was compared to a dipeptidyl peptidase-4 (DPP-4) inhibitor (sitagliptin) and a sulfonylurea (glimepiride).

There have been no clinical efficacy studies conducted with INVOKAMET; however, bioequivalence of INVOKAMET to canagliflozin and metformin co-administered as individual tablets was demonstrated in healthy subjects.

In patients with type 2 diabetes, treatment with canagliflozin and metformin produced clinically and statistically significant improvements in HbA1C compared to placebo. Reductions in HbA1C were observed across subgroups including age, gender, race, and baseline body mass index (BMI).

**INVOKAMET® (canagliflozin and metformin hydrochloride) tablets****14.1 Canagliflozin as Initial Combination Therapy with Metformin**

A total of 1186 patients with type 2 diabetes inadequately controlled with diet and exercise participated in a 26-week double-blind, active-controlled, parallel-group, 5-arm, multicenter study to evaluate the efficacy and safety of initial therapy with canagliflozin in combination with metformin XR. The median age was 56 years, 49% of patients were men, and the mean baseline eGFR was 87.6 mL/min/1.73 m<sup>2</sup>. The median duration of diabetes was 1.6 years, and 72% of patients were treatment naïve. After completing a 2-week single-blind placebo run-in period, patients were randomly assigned for a double-blind treatment period of 26 weeks to 1 of 5 treatment groups (Table 9). The metformin XR dose was initiated at 500 mg/day for the first week of treatment and then increased to 1000 mg/day. Metformin XR or matching placebo was up-titrated every 2-3 weeks during the next 8 weeks of treatment to a maximum daily dose of 1500 to 2000 mg/day, as tolerated; about 90% of patients reached 2000 mg/day.

At the end of treatment, canagliflozin 100 mg and canagliflozin 300 mg in combination with metformin XR resulted in a statistically significant greater improvement in HbA1C compared to their respective canagliflozin doses (100 mg and 300 mg) alone or metformin XR alone.

**Table 9: Results from 26-Week Active-Controlled Clinical Study of Canagliflozin Alone or Canagliflozin as Initial Combination Therapy with Metformin\***

Efficacy Parameter	Metformin XR (N=237)	Canagliflozin 100 mg (N=237)	Canagliflozin 300 mg (N=238)	Canagliflozin 100 mg + Metformin XR (N=237)	Canagliflozin 300 mg + Metformin XR (N=237)
<b>HbA1C (%)</b>					
Baseline (mean)	8.81	8.78	8.77	8.83	8.90
Change from baseline (adjusted mean) <sup>†</sup>	-1.30	-1.37	-1.42	-1.77	-1.78
Difference from canagliflozin 100 mg (adjusted mean) (95% CI) <sup>‡</sup>				-0.40 <sup>‡</sup> (-0.59, -0.21)	
Difference from canagliflozin 300 mg (adjusted mean) (95% CI) <sup>‡</sup>					-0.38 <sup>‡</sup> (-0.56, -0.17)
Difference from metformin XR (adjusted mean) (95% CI) <sup>‡</sup>				-0.46 <sup>‡</sup> (-0.66, -0.27)	-0.46 <sup>‡</sup> (-0.67, -0.26)
Percent of patients achieving HbA1C < 7%	38	34	39	47 <sup>§§</sup>	51 <sup>§§</sup>

\* Intent-to-treat population

<sup>†</sup> Least squares mean adjusted for covariates including baseline value and stratification factor

<sup>‡</sup> Adjusted p<0.001

<sup>§</sup> Adjusted p<0.05

<sup>§§</sup> There were 121 patients without week 26 efficacy data. Analyses addressing missing data gave consistent results with the results provided in this table.

**14.2 Canagliflozin as Add-on Combination Therapy with Metformin**

A total of 1284 patients with type 2 diabetes inadequately controlled on metformin monotherapy (greater than or equal to 2,000 mg/day or at least 1,500 mg/day if higher dose not tolerated) participated in a 26-week, double-blind, placebo- and active-controlled study to evaluate the efficacy and safety of canagliflozin in combination with metformin. The mean age was 55 years, 47% of patients were men, and the mean baseline eGFR was 89 mL/min/1.73 m<sup>2</sup>. Patients already on the required metformin dose (N=1009) were randomized after completing a 2-week, single-blind, placebo run-in period. Patients taking less than the required metformin dose or patients on metformin in combination with another antihyperglycemic agent (N=275) were switched to metformin monotherapy (at doses described above) for at least 8 weeks before entering the 2-week, single-blind, placebo run-in. After the placebo run-in period, patients were randomized to canagliflozin 100 mg, canagliflozin 300 mg, sitagliptin 100 mg, or placebo, administered once daily as add-on therapy to metformin.

At the end of treatment, canagliflozin 100 mg and 300 mg once daily resulted in a statistically significant improvement in HbA1C (p<0.001 for both doses) compared to placebo when added to metformin. Canagliflozin 100 mg and 300 mg once daily

**INVOKAMET® (canagliflozin and metformin hydrochloride) tablets**

also resulted in a greater proportion of patients achieving an HbA1C less than 7%, in significant reduction in fasting plasma glucose (FPG), in improved postprandial glucose (PPG), and in percent body weight reduction compared to placebo when added to metformin (see Table 10). Statistically significant (p<0.001 for both doses) mean changes from baseline in systolic blood pressure relative to placebo were -5.4 mmHg and -6.6 mmHg with canagliflozin 100 mg and 300 mg, respectively.

**Table 10: Results from 26-Week Placebo-Controlled Clinical Study of Canagliflozin in Combination with Metformin\***

Efficacy Parameter	Placebo + Metformin (N=163)	Canagliflozin 100 mg + Metformin (N=368)	Canagliflozin 300 mg + Metformin (N=367)
<b>HbA1C (%)</b>			
Baseline (mean)	7.96	7.94	7.95
Change from baseline (adjusted mean)	-0.17	-0.78	-0.94
Difference from placebo (adjusted mean) (95% CI) <sup>†</sup>		-0.62 <sup>†</sup> (-0.78, -0.48)	-0.77 <sup>†</sup> (-0.91, -0.64)
Percent of patients achieving HbA1C < 7%	30	46 <sup>‡</sup>	58 <sup>‡</sup>
<b>Fasting Plasma Glucose (mg/dL)</b>			
Baseline (mean)	164	169	173
Change from baseline (adjusted mean)	2	-27	-38
Difference from placebo (adjusted mean) (95% CI) <sup>†</sup>		-30 <sup>†</sup> (-36, -24)	-40 <sup>†</sup> (-46, -34)
<b>2-hour Postprandial Glucose (mg/dL)</b>			
Baseline (mean)	249	258	262
Change from baseline (adjusted mean)	-10	-48	-57
Difference from placebo (adjusted mean) (95% CI) <sup>†</sup>		-38 <sup>†</sup> (-49, -27)	-47 <sup>†</sup> (-58, -36)
<b>Body Weight</b>			
Baseline (mean) in kg	88.7	88.7	85.4
% change from baseline (adjusted mean)	-1.2	-3.7	-4.2
Difference from placebo (adjusted mean) (95% CI) <sup>†</sup>		-2.5 <sup>†</sup> (-3.1, -1.9)	-2.9 <sup>†</sup> (-3.5, -2.3)

\* Intent-to-treat population using last observation in study prior to glycemic rescue therapy

<sup>†</sup> Least squares mean adjusted for baseline value and stratification factors

<sup>‡</sup> p<0.001

**14.3 Canagliflozin Compared to Glimepiride, Both as Add-on Combination Therapy with Metformin**

A total of 1450 patients with type 2 diabetes inadequately controlled on metformin monotherapy (greater than or equal to 2,000 mg/day or at least 1,500 mg/day if higher dose not tolerated) participated in a 52-week, double-blind, active-controlled study to evaluate the efficacy and safety of canagliflozin in combination with metformin.

The mean age was 56 years, 52% of patients were men, and the mean baseline eGFR was 90 mL/min/1.73 m<sup>2</sup>. Patients tolerating maximally required metformin dose (N=928) were randomized after completing a 2-week, single-blind, placebo run-in period. Other patients (N=522) were switched to metformin monotherapy (at doses described above) for at least 10 weeks, then completed a 2-week single-blind run-in period. After the 2-week run-in period, patients were randomized to canagliflozin 100 mg, canagliflozin 300 mg, or glimepiride (titration allowed throughout the 52-week study to 8 or 8 mg), administered once daily as add-on therapy to metformin.

As shown in Table 11 and Figure 1, at the end of treatment, canagliflozin 100 mg provided similar reductions in HbA1C from baseline compared to glimepiride when added to metformin therapy. Canagliflozin 300 mg provided a greater reduction from baseline in HbA1C compared to glimepiride, and the relative treatment difference was -0.12% (95% CI: -0.22; -0.02). As shown in Table 11, treatment with canagliflozin 100 mg and 300 mg daily provided greater improvements in percent body weight change, relative to glimepiride.


**INVOKAMET® (canagliflozin and metformin hydrochloride) tablets**

**Table 11: Results from 52-Week Clinical Study Comparing Canagliflozin to Glimepiride in Combination with Metformin\***

Efficacy Parameter	Canagliflozin 100 mg + Metformin (N=483)	Canagliflozin 300 mg + Metformin (N=485)	Glimepiride (Itratred) + Metformin (N=482)
<b>HbA1C (%)</b>			
Baseline (mean)	7.78	7.79	7.83
Change from baseline (adjusted mean)	-0.82	-0.93	-0.81
Difference from glimepiride (adjusted mean) (95% CI)†	-0.01† (-0.11, 0.09)	-0.12† (-0.22, -0.02)	
Percent of patients achieving HbA1C < 7%	54	60	56
<b>Fasting Plasma Glucose (mg/dL)</b>			
Baseline (mean)	165	164	166
Change from baseline (adjusted mean)	-24	-28	-18
Difference from glimepiride (adjusted mean) (95% CI)†	-6 (-10, -2)	-9 (-13, -5)	
<b>Body Weight</b>			
Baseline (mean) in kg	86.8	86.6	86.6
% change from baseline (adjusted mean)	-4.2	-4.7	1.0
Difference from glimepiride (adjusted mean) (95% CI)†	-5.2‡ (-5.7, -4.7)	-5.7‡ (-6.2, -5.1)	

\* Intent-to-treat population using last observation in study prior to glycemic rescue therapy  
 † Least squares mean adjusted for baseline value and stratification factors  
 ‡ Canagliflozin + metformin is considered non-inferior to glimepiride + metformin because the upper limit of this confidence interval is less than the pre-specified non-inferiority margin of < 0.3%.  
 § p<0.001

**Figure 1: Mean HbA1C Changes at Each Time Point (Completers) and at Week 52 Using Last Observation Carried Forward (mITT Population)**



**14.4 Canagliflozin as Add-on Combination Therapy with Metformin and Sulfonyleurea**

A total of 469 patients with type 2 diabetes inadequately controlled on the combination of metformin (greater than or equal to 2,000 mg/day or at least 1,500 mg/day if higher dose not tolerated) and sulfonyleurea (maximal or near-maximal effective dose) participated in a 26-week, double-blind, placebo-controlled study to evaluate the efficacy and safety of canagliflozin in combination with metformin and sulfonyleurea. The mean age was 57 years, 51% of patients were men, and the mean baseline eGFR was 89 mL/min/1.73 m<sup>2</sup>. Patients already on the protocol-specified doses of metformin and sulfonyleurea (N=372) entered a 2-week, single-blind, placebo run-in period. Other patients (N=97) were required to be on a stable protocol-specified dose of metformin and sulfonyleurea

**INVOKAMET® (canagliflozin and metformin hydrochloride) tablets**

for at least 8 weeks before entering the 2-week run-in period. Following the run-in period, patients were randomized to canagliflozin 100 mg, canagliflozin 300 mg, or placebo administered once daily as add-on to metformin and sulfonyleurea.

At the end of treatment, canagliflozin 100 mg and 300 mg once daily resulted in a statistically significant improvement in HbA1C (p<0.001 for both doses) compared to placebo when added to metformin and sulfonyleurea. Canagliflozin 100 mg and 300 mg once daily also resulted in a greater proportion of patients achieving an HbA1C less than 7.0%, in a significant reduction in fasting plasma glucose (FPG), and in percent body weight reduction compared to placebo when added to metformin and sulfonyleurea (see Table 12).

**Table 12: Results from 26-Week Placebo-Controlled Clinical Study of Canagliflozin in Combination with Metformin and Sulfonyleurea\***

Efficacy Parameter	Placebo + Metformin and Sulfonyleurea (N=156)	Canagliflozin 100 mg + Metformin and Sulfonyleurea (N=157)	Canagliflozin 300 mg + Metformin and Sulfonyleurea (N=156)
<b>HbA1C (%)</b>			
Baseline (mean)	8.12	8.13	8.13
Change from baseline (adjusted mean)	-0.13	-0.85	-1.06
Difference from placebo (adjusted mean) (95% CI)†		-0.71‡ (-0.80, -0.52)	-0.92‡ (-1.11, -0.73)
Percent of patients achieving HbA1C < 7%	18	43‡	57‡
<b>Fasting Plasma Glucose (mg/dL)</b>			
Baseline (mean)	170	173	169
Change from baseline (adjusted mean)	4	-18	-31
Difference from placebo (adjusted mean) (95% CI)†		-22‡ (-31, -13)	-35† (-44, -25)
<b>Body Weight</b>			
Baseline (mean) in kg	90.8	93.5	93.5
% change from baseline (adjusted mean)	-0.7	-2.1	-2.6
Difference from placebo (adjusted mean) (95% CI)†		-1.4‡ (-2.1, -0.7)	-2.0‡ (-2.7, -1.3)

\* Intent-to-treat population using last observation in study prior to glycemic rescue therapy  
 † Least squares mean adjusted for baseline value and stratification factors  
 ‡ p<0.001

**14.5 Canagliflozin Compared to Sitagliptin, Both as Add-on Combination Therapy with Metformin and Sulfonyleurea**

A total of 755 patients with type 2 diabetes inadequately controlled on the combination of metformin (greater than or equal to 2,000 mg/day or at least 1,500 mg/day if higher dose not tolerated) and sulfonyleurea (near-maximal or maximal effective dose) participated in a 52 week, double-blind, active-controlled study to compare the efficacy and safety of canagliflozin 300 mg versus sitagliptin 100 mg in combination with metformin and sulfonyleurea. The mean age was 57 years, 56% of patients were men, and the mean baseline eGFR was 88 mL/min/1.73 m<sup>2</sup>. Patients already on protocol-specified doses of metformin and sulfonyleurea (N=716) entered a 2-week single-blind, placebo run-in period. Other patients (N=39) were required to be on a stable protocol-specified dose of metformin and sulfonyleurea for at least 8 weeks before entering the 2-week run-in period. Following the run-in period, patients were randomized to canagliflozin 300 mg or sitagliptin 100 mg as add-on to metformin and sulfonyleurea.

As shown in Table 13 and Figure 2, at the end of treatment, canagliflozin 300 mg provided greater HbA1C reduction compared to sitagliptin 100 mg when added to metformin and sulfonyleurea (p<0.05). Canagliflozin 300 mg resulted in a mean percent change in body weight from baseline of -2.5% compared to +0.3% with sitagliptin 100 mg. A mean change in systolic blood pressure from baseline of -5.06 mmHg was observed with canagliflozin 300 mg compared to +0.85 mmHg with sitagliptin 100 mg.

**INVOKAMET® (canagliflozin and metformin hydrochloride) tablets**

**Table 13: Results from 52-Week Clinical Study Comparing Canagliflozin to Sitagliptin in Combination with Metformin and Sulfonylurea\***

Efficacy Parameter	Canagliflozin 300 mg + Metformin and Sulfonylurea (N=377)	Sitagliptin 100 mg + Metformin and Sulfonylurea (N=378)
<b>HbA1C (%)</b>		
Baseline (mean)	8.12	8.13
Change from baseline (adjusted mean)	-1.03	-0.66
Difference from sitagliptin (adjusted mean) (95% CI) <sup>†</sup>	-0.37 <sup>‡</sup>	(-0.50, -0.25)
Percent of patients achieving HbA1C < 7%	48	35
<b>Fasting Plasma Glucose (mg/dL)</b>		
Baseline (mean)	170	164
Change from baseline (adjusted mean)	-30	-6
Difference from sitagliptin (adjusted mean) (95% CI) <sup>†</sup>	-24	(-30, -18)
<b>Body Weight</b>		
Baseline (mean) in kg	87.6	89.6
% change from baseline (adjusted mean)	-2.5	0.3
Difference from sitagliptin (adjusted mean) (95% CI) <sup>†</sup>	-2.8 <sup>‡</sup>	(-3.3, -2.2)


\* Intent-to-treat population using last observation in study prior to glycemic rescue therapy

<sup>†</sup> Least squares mean adjusted for baseline value and stratification factors

<sup>‡</sup> Canagliflozin + metformin + sulfonylurea is considered noninferior to sitagliptin + metformin + sulfonylurea because the upper limit of this confidence interval is less than the pre-specified non-inferiority margin of < 0.3%.

<sup>§</sup> p<0.001

**Figure 2: Mean HbA1C Change at Each Time Point (Completers) and at Week 52 Using Last Observation Carried Forward (mITT Population)**



**14.6 Canagliflozin as Add-on Combination Therapy with Metformin and Pioglitazone**

A total of 342 patients with type 2 diabetes inadequately controlled on the combination of metformin (greater than or equal to 2,000 mg/day or at least 1,500 mg/day if higher dose not tolerated) and pioglitazone (30 or 45 mg/day) participated in a 26-week, double-blind, placebo-controlled study to evaluate the efficacy and safety of canagliflozin in combination with metformin and pioglitazone. The mean age was 57 years, 63% of patients were men, and the mean baseline eGFR was 66 mL/min/1.73 m<sup>2</sup>. Patients already on protocol-specified doses of metformin and pioglitazone (N=163) entered a 2-week, single-blind, placebo run-in period. Other patients (N=181) were required to be on stable protocol-specified doses of metformin and pioglitazone for at least 8 weeks before entering the 2-week run-in period. Following the run-in period, patients were randomized to canagliflozin 100 mg, canagliflozin 300 mg, or placebo, administered once daily as add-on to metformin and pioglitazone.

At the end of treatment, canagliflozin 100 mg and 300 mg once daily resulted in a statistically significant improvement in HbA1C (p<0.001 for both doses) compared to placebo when added to metformin and pioglitazone. Canagliflozin 100 mg and 300 mg once daily also resulted in a greater proportion of patients

**INVOKAMET® (canagliflozin and metformin hydrochloride) tablets**

achieving an HbA1C less than 7%, in significant reduction in fasting plasma glucose (FPG), and in percent body weight reduction compared to placebo when added to metformin and pioglitazone (see Table 14). Statistically significant (p<0.05 for both doses) mean changes from baseline in systolic blood pressure relative to placebo were -4.1 mmHg and -3.5 mmHg with canagliflozin 100 mg and 300 mg, respectively.

**Table 14: Results from 26-Week Placebo-Controlled Clinical Study of Canagliflozin in Combination with Metformin and Pioglitazone\***

Efficacy Parameter	Placebo + Metformin and Pioglitazone (N=115)	Canagliflozin 100 mg + Metformin and Pioglitazone (N=113)	Canagliflozin 300 mg + Metformin and Pioglitazone (N=114)
<b>HbA1C (%)</b>			
Baseline (mean)	8.00	7.99	7.84
Change from baseline (adjusted mean)	-0.26	-0.89	-1.03
Difference from placebo (adjusted mean) (95% CI) <sup>†</sup>		-0.62 <sup>‡</sup>	-0.76 <sup>‡</sup>
		(-0.81, -0.44)	(-0.85, -0.58)
Percent of patients achieving HbA1C < 7%	33	47 <sup>‡</sup>	64 <sup>‡</sup>
<b>Fasting Plasma Glucose (mg/dL)</b>			
Baseline (mean)	164	169	164
Change from baseline (adjusted mean)	3	-27	-33
Difference from placebo (adjusted mean) (95% CI) <sup>†</sup>		-29 <sup>‡</sup>	-36 <sup>‡</sup>
		(-37, -22)	(-43, -28)
<b>Body Weight</b>			
Baseline (mean) in kg	94.0	94.2	94.4
% change from baseline (adjusted mean)	-0.1	-2.8	-3.8
Difference from placebo (adjusted mean) (95% CI) <sup>†</sup>		-2.7 <sup>‡</sup>	-3.7 <sup>‡</sup>
		(-3.6, -1.8)	(-4.6, -2.8)

\* Intent-to-treat population using last observation in study prior to glycemic rescue therapy

<sup>†</sup> Least squares mean adjusted for baseline value and stratification factors

<sup>‡</sup> p<0.001

**14.7 Canagliflozin as Add-on Combination Therapy with Insulin (With or Without Other Anti-Hyperglycemic Agents, Including Metformin)**

A total of 1718 patients with type 2 diabetes inadequately controlled on insulin greater than or equal to 30 units/day or insulin in combination with other antihyperglycemic agents participated in an 18-week, double-blind, placebo-controlled substudy of a cardiovascular study to evaluate the efficacy and safety of canagliflozin in combination with insulin. Of these patients, a subgroup of 432 patients with inadequate glycemic control received canagliflozin or placebo plus metformin and ≥ 30 units/day of insulin over 18 weeks.

In this subgroup, the mean age was 61 years, 67% of patients were men, and the mean baseline eGFR was 81 mL/min/1.73 m<sup>2</sup>. Patients on metformin in combination with basal, bolus, or basal/bolus insulin for at least 10 weeks entered a 2-week, single-blind, placebo run-in period. Approximately 74% of these patients were on a background of metformin and basal/bolus insulin regimen. After the run-in period, patients were randomized to canagliflozin 100 mg, canagliflozin 300 mg, or placebo, administered once daily as add-on to metformin and insulin. The mean daily insulin dose at baseline was 93 units, which was similar across treatment groups.

At the end of treatment, canagliflozin 100 mg and 300 mg once daily resulted in a statistically significant improvement in HbA1C (p<0.001 for both doses) compared to placebo when added to metformin and insulin. Canagliflozin 100 mg and 300 mg once daily also resulted in a greater proportion of patients achieving an HbA1C less than 7%, in significant reductions in fasting plasma glucose (FPG), and in percent body weight reductions compared to placebo (see Table 15). Statistically significant (p=0.023 for the 100 mg and p<0.001 for the 300 mg dose) mean change from baseline in systolic blood pressure relative to placebo was -3.5 mmHg and -6 mmHg with canagliflozin 100 mg and 300 mg, respectively. Fewer patients on canagliflozin in combination with metformin and insulin required glycemic rescue therapy: 3.6% of patients receiving canagliflozin 100 mg, 2.7% of patients receiving canagliflozin 300 mg, and 6.2% of patients receiving placebo. An increased incidence of hypoglycemia was observed in this study, which is consistent with the expected increase of hypoglycemia when an agent not associated with hypoglycemia is added to insulin (see Warnings and Precautions (5.7), Adverse Reactions (6.1)).

**INVOKAMET® (canagliflozin and metformin hydrochloride) tablets****Table 15: Results from 18-Week Placebo-Controlled Clinical Study of Canagliflozin in Combination with Metformin and Insulin  $\geq 30$  Units/Day\***

Efficacy Parameter	Placebo + Metformin + Insulin (N=145)	Canagliflozin 100 mg + Metformin + Insulin (N=139)	Canagliflozin 300 mg + Metformin + Insulin (N=146)
<b>HbA1C (%)</b>			
Baseline (mean)	8.15	8.20	8.22
Change from baseline (adjusted mean)	0.03	-0.64	-0.79
Difference from placebo (adjusted mean) (95% CI) <sup>†</sup>		-0.66 <sup>‡</sup> (-0.81, -0.51)	-0.82 <sup>‡</sup> (-0.98, -0.67)
Percent of patients achieving HbA1C < 7%	9	19 <sup>§</sup>	29 <sup>§</sup>
<b>Fasting Plasma Glucose (mg/dL)</b>			
Baseline	163	168	167
Change from baseline (adjusted mean)	1	-16	-24
Difference from placebo (adjusted mean) (97.5% CI) <sup>†</sup>		-16 <sup>‡</sup> (-28, -5)	-25 <sup>‡</sup> (-36, -14)
<b>Body Weight</b>			
Baseline (mean) in kg	102.3	99.7	101.1
% change from baseline (adjusted mean)	0.0	-1.7	-2.7
Difference from placebo (adjusted mean) (97.5% CI) <sup>†</sup>		-1.7 <sup>‡</sup> (-2.4, -1.0)	-2.7 <sup>‡</sup> (-3.4, -2.0)

\* Intent-to-treat population using last observation in study prior to glycemic rescue therapy

<sup>†</sup> Least squares mean adjusted for baseline value and stratification factors

<sup>‡</sup>  $p \leq 0.001$

<sup>§</sup>  $p \leq 0.01$

**16 HOW SUPPLIED/STORAGE AND HANDLING**

INVOKAMET (canagliflozin and metformin hydrochloride) tablets are available in the strengths and packages listed below:

Canagliflozin 50 mg and metformin hydrochloride 500 mg tablets are immediate-release, capsule-shaped, white film-coated tablets with "CM" on one side and "155" on the other side.

• NDC 50458-540-60 Bottle of 60

Canagliflozin 50 mg and metformin hydrochloride 1,000 mg tablets are immediate-release, capsule-shaped, beige film-coated tablets with "CM" on one side and "551" on the other side.

• NDC 50458-541-60 Bottle of 60

Canagliflozin 150 mg and metformin hydrochloride 500 mg tablets are immediate-release, capsule-shaped, yellow film-coated tablets with "CM" on one side and "215" on the other side.

• NDC 50458-542-60 Bottle of 60

Canagliflozin 150 mg and metformin hydrochloride 1,000 mg tablets are immediate-release, capsule-shaped, purple film-coated tablets with "CM" on one side and "611" on the other side.

• NDC 50458-543-60 Bottle of 60

**Storage and Handling**

Keep out of reach of children.

Store at 68-77°F (20-25°C); excursions permitted between 59°F and 86°F (15°C and 30°C) [see USP Controlled Room Temperature]. Store in the original container.

**17 PATIENT COUNSELING INFORMATION**

Advise the patient to read the FDA-Approved Patient Labeling (Medication Guide).

- **Lactic Acidosis:** Explain the risks of lactic acidosis, its symptoms, and conditions that predispose to its development, as noted in *Warnings and Precautions (5.1)*. Advise patients to discontinue INVOKAMET immediately and to promptly notify their healthcare provider if unexplained hyperventilation, myalgias, malaise, unusual somnolence or other nonspecific symptoms occur. Once a patient is stabilized on INVOKAMET, gastrointestinal symptoms, which are common during initiation of metformin, are unlikely to recur. Later occurrence of gastrointestinal symptoms could be due to lactic acidosis or other serious disease.
- Instruct patients to keep INVOKAMET in the original bottle to protect from moisture. Do not put INVOKAMET in pill boxes or pill organizers.
- Counsel patients against excessive alcohol intake while receiving INVOKAMET.
- Inform patients about importance of regular testing of renal function and hematological parameters while receiving INVOKAMET.

**INVOKAMET® (canagliflozin and metformin hydrochloride) tablets**

- Advise patients to seek medical advice promptly during periods of stress such as fever, trauma, infection, or surgery, as medication requirements may change.
- Instruct patients to take INVOKAMET only as prescribed twice daily with food. If a dose is missed, advise patients not to take two doses of INVOKAMET at the same time.
- **Hypotension:** Inform patients that symptomatic hypotension may occur with INVOKAMET and advise them to contact their doctor if they experience such symptoms [see *Warnings and Precautions (5.2)*]. Inform patients that dehydration may increase the risk for hypotension and to have adequate fluid intake.
- **Ketoacidosis:** Inform patients that ketoacidosis is a serious life-threatening condition. Cases of ketoacidosis have been reported during use of canagliflozin. Instruct patients to check ketones (when possible) if symptoms consistent with ketoacidosis occur even if blood glucose is not elevated. If symptoms of ketoacidosis (including nausea, vomiting, abdominal pain, tiredness, and labored breathing) occur, instruct patients to discontinue INVOKAMET and seek medical advice immediately [see *Warnings and Precautions (5.3)*].
- **Acute Kidney Injury:** Inform patients that acute kidney injury has been reported during use of canagliflozin. Advise patients to seek medical advice immediately if they have reduced oral intake (such as due to acute illness or fasting), or increased fluid losses (such as due to vomiting, diarrhea, or excessive heat exposure), as it may be appropriate to temporarily discontinue INVOKAMET use in those settings [see *Warnings and Precautions (5.4)*].
- **Serious Urinary Tract Infections:** Inform patients of the potential for urinary tract infections, which may be serious. Provide them with information on the symptoms of urinary tract infections. Advise them to seek medical advice if such symptoms occur [see *Warnings and Precautions (5.6)*].
- **Genital Mycotic Infections in Females:** Inform female patients that vaginal yeast infection (e.g., vulvovaginitis) may occur and provide them with information on the signs and symptoms of a vaginal yeast infection. Advise them of treatment options and when to seek medical advice [see *Warnings and Precautions (5.8)*].
- **Genital Mycotic Infections in Males:** Inform male patients that yeast infection of the penis (e.g., balanitis or balanoposthitis) may occur, especially in uncircumcised males and patients with prior history. Provide them with information on the signs and symptoms of balanitis and balanoposthitis (rash or redness of the glans or foreskin of the penis). Advise them of treatment options and when to seek medical advice [see *Warnings and Precautions (5.8)*].
- **Hypersensitivity Reactions:** Inform patients that serious hypersensitivity reactions, such as urticaria, rash, anaphylaxis, and angioedema, have been reported with canagliflozin. Advise patients to report immediately any signs or symptoms suggesting allergic reaction and to discontinue drug until they have consulted prescribing physicians [see *Warnings and Precautions (5.9)*].
- **Bone Fracture:** Inform patients that bone fractures have been reported in patients taking canagliflozin. Provide them with information on factors that may contribute to fracture risk.
- **Laboratory Tests:** Inform patients that they will test positive for glucose in their urine while on INVOKAMET [see *Drug Interactions (7.2)*].
- **Pregnancy:** Advise pregnant women, and females of reproductive potential of the potential risk to a fetus with treatment with INVOKAMET [see *Use in Specific Populations (8.1)*]. Instruct females of reproductive potential to report pregnancies to their physicians as soon as possible.
- **Lactation:** Advise women that breastfeeding is not recommended during treatment with INVOKAMET [see *Use in Specific Populations (8.2)*].
- Inform females that treatment with INVOKAMET may result in ovulation in some premenopausal anovulatory women which may lead to unintended pregnancy [see *Use in Specific Populations (8.3)*].
- Inform patients that the most common adverse reactions associated with canagliflozin are genital mycotic infection, urinary tract infection, and increased urination. Most common adverse reactions associated with metformin are diarrhea, nausea, vomiting, flatulence, asthenia, indigestion, abdominal discomfort, and headache.

Manufactured by:  
Janssen Pharmaceuticals, Inc.  
Titusville, NJ 08580

Finished product manufactured by:  
Janssen Ortho, LLC  
Gurabo, PR 00778

Licensed from Mitsubishi Tanabe Pharma Corporation



© 2014 Janssen Pharmaceuticals, Inc.

058065-160811

**Medication Guide**  
**INVOKAMET® (in vok' a met)**  
**(canagliflozin and metformin hydrochloride)**  
**Tablets**

**What is the most important information I should know about INVOKAMET?**

**INVOKAMET can cause serious side effects, including:**

- **Lactic Acidosis.** Metformin, one of the medicines in INVOKAMET, can cause a rare but serious condition called lactic acidosis (a build-up of lactic acid in the blood) that can cause death. Lactic acidosis is a medical emergency and must be treated in the hospital.

**Call your doctor right away if you have any of the following symptoms, which could be signs of lactic acidosis:**

- you feel cold in your hands or feet
- you feel very weak or tired
- you have trouble breathing
- you have stomach pains, nausea, or vomiting
- you have a slow or irregular heartbeat
- you have unusual (not normal) muscle pain
- you have unusual sleepiness or sleep longer than usual
- you feel dizzy or lightheaded

Most people who have had lactic acidosis had other conditions that, in combination with metformin use, led to the lactic acidosis. Tell your doctor if you have any of the following, because you have a higher chance for getting lactic acidosis with INVOKAMET if you:

- have severe kidney problems or your kidneys are affected by certain x-ray tests that use injectable dye.
- have liver problems
- drink alcohol very often, or drink a lot of alcohol in short-term "binge" drinking.
- get dehydrated (lose a large amount of body fluids). This can happen if you are sick with a fever, vomiting, or diarrhea. Dehydration can also happen when you sweat a lot with activity or exercise and do not drink enough fluids.
- have surgery.
- have a heart attack, severe infection, or stroke.

The best way to keep from having a problem with lactic acidosis from metformin is to tell your doctor if you have any of the problems in the list above. Your doctor will decide to stop your INVOKAMET for a while if you have any of these things.

**INVOKAMET can have other serious side effects. See "What are the possible side effects of INVOKAMET?"**

**What is INVOKAMET?**

- INVOKAMET contains 2 prescription medicines called canagliflozin (INVOKANA) and metformin hydrochloride (GLUCOPHAGE). INVOKAMET can be used along with diet and exercise to improve blood sugar (glucose) control in adults with type 2 diabetes when treatment with both canagliflozin and metformin is appropriate.
- INVOKAMET is not for people with diabetic ketoacidosis (increased ketones in blood or urine).
- It is not known if INVOKAMET is safe and effective in children under 18 years of age.

**Who should not take INVOKAMET?**

**Do not take INVOKAMET if you:**

- have moderate to severe kidney problems or are on dialysis.
- have a condition called metabolic acidosis or diabetic ketoacidosis (increased ketones in the blood or urine).
- are allergic to canagliflozin, metformin, or any of the ingredients in INVOKAMET. See the end of this Medication Guide for a list of ingredients in INVOKAMET. Symptoms of allergic reaction to INVOKAMET may include:
  - rash
  - raised red patches on your skin (hives)
  - swelling of the face, lips, mouth, tongue, and throat that may cause difficulty in breathing or swallowing

**INVOKAMET® (canagliflozin and metformin hydrochloride) tablets****What should I tell my doctor before taking INVOKAMET?**

Before you take INVOKAMET, tell your doctor if you:

- have moderate to severe kidney problems.
- have liver problems.
- have a history of urinary tract infections or problems with urination.
- are on a low sodium (salt) diet. Your doctor may change your diet or your dose of INVOKAMET.
- have ever had an allergic reaction to INVOKAMET.
- are going to get an injection of dye or contrast agents for an x-ray procedure. INVOKAMET may need to be stopped for a short time. Talk to your doctor about when you should stop INVOKAMET and when you should start INVOKAMET again. See **"What is the most important information I should know about INVOKAMET?"**
- have heart problems, including congestive heart failure.
- are going to have surgery.
- are eating less due to illness, surgery, or a change in your diet.
- have or have had problems with your pancreas, including pancreatitis or surgery on your pancreas.
- drink alcohol very often, or drink a lot of alcohol in the short-term ("binge" drinking).
- have other medical conditions.
- are pregnant or plan to become pregnant. INVOKAMET may harm your unborn baby. If you become pregnant while taking INVOKAMET, tell your doctor as soon as possible. Talk with your doctor about the best way to control your blood sugar while you are pregnant.
- are a premenopausal woman (before the "change of life"), who does not have periods regularly or at all. INVOKAMET may increase your chance of becoming pregnant. Talk to your doctor about birth control choices while taking INVOKAMET, if you are not planning to become pregnant. Tell your doctor right away if you become pregnant while taking INVOKAMET.
- are breastfeeding or plan to breastfeed. INVOKAMET may pass into your breast milk and may harm your baby. Talk with your doctor about the best way to feed your baby if you are taking INVOKAMET. Do not breastfeed while taking INVOKAMET.

**Tell your doctor about all the medicines you take, including prescription and over-the-counter medicines, vitamins, and herbal supplements.**

INVOKAMET may affect the way other medicines work and other medicines may affect how INVOKAMET works. Especially tell your doctor if you take:

- |   |  |
|---|--|
| • diuretics (water pills)                               | • rifampin (used to treat or prevent tuberculosis)             |
| • phenytoin or phenobarbital (used to control seizures) | • ritonavir (Norvir®, Kaletra®)* (used to treat HIV infection) |
| • digoxin (Lanoxin®)* (used to treat heart problems)    |  |

Ask your doctor or pharmacist for a list of these medicines if you are not sure if your medicine is listed above.

Know the medicines you take. Keep a list of them and show it to your doctor and pharmacist when you get a new medicine.

**How should I take INVOKAMET?**

- Take INVOKAMET by mouth 2 times each day with meals exactly as your doctor tells you to take it. Taking INVOKAMET with meals may lower your chance of having an upset stomach.
- Your doctor will tell you how much INVOKAMET to take and when to take it. Your doctor may change your dose if needed.
- Your doctor may tell you to take INVOKAMET along with other diabetes medicines. Low blood sugar can happen more often when INVOKAMET is taken with certain other diabetes medicines. See **"What are the possible side effects of INVOKAMET?"**
- If you miss a dose, take it as soon as you remember. If it is almost time for your next dose, skip the missed dose and take the medicine at the next regularly scheduled time. Do not take 2 doses of INVOKAMET at the same time. Talk to your doctor if you have questions about a missed dose.
- If you take too much INVOKAMET, call your doctor or go to the nearest hospital emergency room right away.
- When your body is under some types of stress, such as fever, trauma (such as a car accident), infection, or surgery, the amount of diabetes medicine you need may change. Tell your doctor right away if you have any of these conditions and follow your doctor's instructions.
- Stay on your prescribed diet and exercise program while taking INVOKAMET.
- Check your blood sugar as your doctor tells you to.
- INVOKAMET will cause your urine to test positive for glucose.
- Your doctor may do certain blood tests before you start INVOKAMET and during treatment as needed. Your doctor may change your dose of INVOKAMET based on the results of your blood tests.
- Your doctor will check your diabetes with regular blood tests, including your blood sugar levels and your hemoglobin A1C.

**INVOKAMET® (canagliflozin and metformin hydrochloride) tablets****What should I avoid while taking INVOKAMET?**

- Avoid drinking alcohol very often, or drinking a lot of alcohol in a short period of time ("binge" drinking). It can increase your chances of getting serious side effects.

**What are the possible side effects of INVOKAMET?****INVOKAMET may cause serious side effects including:**

- See "What is the most important information I should know about INVOKAMET?"
- **dehydration.** INVOKAMET can cause some people to become dehydrated (the loss of too much body water). Dehydration may cause you to feel dizzy, faint, lightheaded, or weak, especially when you stand up (orthostatic hypotension).

You may be at higher risk of dehydration if you:

- have low blood pressure
- take medicines to lower your blood pressure, including diuretics (water pill)
- are on a low sodium (salt) diet
- have kidney problems
- are 65 years of age or older

Talk to your doctor about what you can do to prevent dehydration including how much fluid you should drink on a daily basis.

- **ketoacidosis (Increased ketones in your blood or urine).** Ketoacidosis has happened in people who have **type 1 diabetes or type 2 diabetes**, during treatment with canagliflozin, one of the medicines in INVOKAMET. Ketoacidosis is a serious condition, which may need to be treated in a hospital. Ketoacidosis may lead to death. **Ketoacidosis can happen with INVOKAMET, even if your blood sugar is less than 250 mg/dL. Stop taking INVOKAMET and call your doctor right away if you get any of the following symptoms:**

- nausea
- vomiting
- stomach area (abdominal) pain
- tiredness
- trouble breathing

If you get any of these symptoms during treatment with INVOKAMET, if possible, check for ketones in your urine, even if your blood sugar is less than 250 mg/dL.

- **kidney problems.** Sudden kidney injury has happened to people taking INVOKAMET. Talk to your doctor right away if you:
  - reduce the amount of food or liquid you drink for example, if you are sick or cannot eat or
  - you start to lose liquids from your body for example, from vomiting, diarrhea or being in the sun too long.
- a high amount of potassium in your blood.
- **serious urinary tract infections.** Serious urinary tract infections that may lead to hospitalization have happened in people who are taking canagliflozin, one of the medicines in INVOKAMET. Tell your doctor if you have any signs or symptoms of a urinary tract infection such as a burning feeling when passing urine, a need to urinate often, the need to urinate right away, pain in the lower part of your stomach (pelvis), or blood in the urine. Sometimes people may also have a fever, back pain, nausea, or vomiting.
- **low blood sugar (hypoglycemia).** If you take INVOKAMET with another medicine that can cause low blood sugar, such as a sulfonylurea or insulin, your risk of getting low blood sugar is higher. The dose of your sulfonylurea medicine or insulin may need to be lowered while you take INVOKAMET. Signs and symptoms of low blood sugar may include:
 

◦ headache	◦ drowsiness	◦ weakness	◦ confusion	◦ dizziness
◦ irritability	◦ hunger	◦ fast heartbeat	◦ sweating	◦ shaking or feeling jittery
- **vaginal yeast infection.** Women who take INVOKAMET may get vaginal yeast infections. Symptoms of a vaginal yeast infection include:
  - vaginal odor
  - white or yellowish vaginal discharge (discharge may be lumpy or look like cottage cheese)
  - vaginal itching
- **yeast infection of the penis (balanitis or balanoposthitis).** Men who take INVOKAMET may get a yeast infection of the skin around the penis. Certain men who are not circumcised may have swelling of the penis that makes it difficult to pull back the skin around the tip of the penis. Other symptoms of yeast infection of the penis include:
 

◦ redness, itching, or swelling of the penis	◦ rash of the penis
◦ foul smelling discharge from the penis	◦ pain in the skin around the penis

**INVOKAMET® (canagliflozin and metformin hydrochloride) tablets****What are the possible side effects of INVOKAMET? (continued)**

Talk to your doctor about what to do if you get symptoms of a yeast infection of the vagina or penis. Your doctor may suggest you use an over-the-counter antifungal medicine. Talk to your doctor right away if you use an over-the-counter antifungal medication and your symptoms do not go away.

- **serious allergic reaction.** If you have any symptoms of a serious allergic reaction, stop taking INVOKAMET and call your doctor right away or go to the nearest hospital emergency room. See **"Who should not take INVOKAMET?"**. Your doctor may give you a medicine for your allergic reaction and prescribe a different medicine for your diabetes.
- **broken bones (fractures).** Bone fractures have been seen in patients taking canagliflozin. Talk to your doctor about factors that may increase your risk of bone fracture.
- **low vitamin B<sub>12</sub> (vitamin B<sub>12</sub> deficiency).** Using metformin for long periods of time may cause a decrease in the amount of vitamin B<sub>12</sub> in your blood, especially if you have had low vitamin B<sub>12</sub> blood levels before. Your doctor may do blood tests to check your vitamin B<sub>12</sub> levels.

Other common side effects of INVOKAMET include:

- nausea and vomiting
- diarrhea
- weakness
- gas
- upset stomach
- indigestion
- headache
- changes in urination, including urgent need to urinate more often, in larger amounts, or at night

Tell your doctor if you have any side effect that bothers you or that does not go away. These are not all the possible side effects of INVOKAMET. For more information, ask your doctor or pharmacist.

**Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.**

You may also report side effects to Janssen Pharmaceuticals, Inc. at 1-800-526-7736.

**How should I store INVOKAMET?**

- Store INVOKAMET at room temperature between 68°F to 77°F (20°C to 25°C).
- Store in the original container to protect from moisture. **Do not put INVOKAMET in pill boxes or pill organizers.**

**Keep INVOKAMET and all medicines out of the reach of children.**

**General information about the safe and effective use of INVOKAMET.**

Medicines are sometimes prescribed for purposes other than those listed in the Medication Guide. Do not use INVOKAMET for a condition for which it was not prescribed. Do not give INVOKAMET to other people, even if they have the same symptoms you have. It may harm them.

This Medication Guide summarizes the most important information about INVOKAMET. If you would like more information, talk with your doctor. You can ask your pharmacist or doctor for information about INVOKAMET that is written for healthcare professionals.

**For more information about INVOKAMET, call 1-800-526-7736 or visit our website at [www.invokamet.com](http://www.invokamet.com).**

**What are the ingredients of INVOKAMET?**

Active ingredients: canagliflozin and metformin hydrochloride

Inactive ingredients: The tablet core contains croscarmellose sodium, hypromellose, magnesium stearate, and microcrystalline cellulose. The magnesium stearate is vegetable-sourced. In addition, the tablet coating contains Macrogol/PEG, polyvinyl alcohol (partially hydrolyzed), talc, titanium dioxide, iron oxide yellow (50 mg/1,000 mg and 150 mg/500 mg tablets only), iron oxide red (50 mg/1,000 mg, 150 mg/500 mg and 150 mg/1,000 mg tablets only), and iron oxide black (150 mg/1,000 mg tablets only).

\* The brands listed are trademarks of their respective owners and are not trademarks of Janssen Pharmaceuticals, Inc.

Manufactured for: Janssen Pharmaceuticals, Inc., Titusville, NJ 08560. Manufactured by: Janssen Ortho, LLC, Gurabo, PR 00778. Licensed from Mitsubishi Tanabe Pharma Corporation. © 2014 Janssen Pharmaceuticals, Inc.

This Medication Guide has been approved by the U.S. Food and Drug Administration.

Revised 07/2016

058065-160811