

Neocate® Mixing Instructions



Neocate Infant DHA / ARA



Standard Dilution 20 kcal/fl oz	Amount of Water to Add		Amount of Neocate Powder‡	Final Volume	
	mL	fl oz†		mL	fl oz†
	90	3	3 scoops (13.8 g)	100	3.3
	150	5	5 scoops (23 g)	166	5.5
	240	8	8 scoops (37 g)	266	9
	480	16	16 scoops (73.6 g)	531	17.7
	720	24	24 scoops (110.4 g)	797	26.6

Neocate Infant DHA / ARA Mixing Instructions

Neocate Infant DHA/ARA may be mixed with warm or cool sterile water. Do not boil formula or use microwave to prepare or warm formula. For standard dilution, use enclosed scoop (4.6 g) to mix 1 level, unpacked scoop to every fluid ounce (fl oz) of water. Example: To make ~4 fl oz: 1) Add sterile water to a clean, sterile container up to the 4 fl oz line. 2) Add 4 leveled and unpacked scoops of Neocate Infant powder. 3) Close and shake for 15-20 seconds or until powder is dissolved. Freshly prepared formula is best, but formula prepared in advance may be cooled rapidly and stored in refrigerator and used within 24 hours. Shake before use. Formula may be kept at room temperature for 4 hours. Once feeding begins, use within 1 hour. May be used for oral or tube feeding.

Storage: Do not store at extreme temperatures. Cover and store opened cans in a cool, dry place (not a refrigerator). Use within one month after opening.

Neocate Junior, Chocolate & Tropical



Standard Dilution 30 kcal/fl oz	Amount of Water to Add		Amount of Neocate Powder‡	Final Volume	
	mL	fl oz†		mL	fl oz†
	124	4	4 scoops (32.8 g)	148	5
	218	7	7 scoops (57.4 g)	259	8
	404	13.5	13 scoops (107 g)	481	16
	622	20	20 scoops (164 g)	740	24
	808	27	26 scoops (213.2 g)	962	32

Neocate Junior Unflavored & Neocate Junior with Prebiotics, Unflavored



Standard Dilution 30 kcal/fl oz	Amount of Water to Add		Amount of Neocate Powder‡	Final Volume	
	mL	fl oz†		mL	fl oz†
	100	3.5	3 scoops (24.6 g)	118	4
	199	6.5	6 scoops (49.2 g)	235	8
	299	10	9 scoops (73.8 g)	353	12
	598	20	18 scoops (147.6 g)	706	24
	814	27	24 scoops (196.8 g)	950	32

Neocate Junior with Prebiotics, Vanilla



Standard Dilution 30 kcal/fl oz	Amount of Water to Add		Amount of Neocate Powder‡	Final Volume	
	mL	fl oz†		mL	fl oz†
	127	4	4 scoops (32.8 g)	151	5
	223	7.5	7 scoops (57.4 g)	263	9
	382	13	12 scoops (98.4 g)	452	15
	636	21	20 scoops (164 g)	753	25
	891	29.5	28 scoops (229.6 g)	1054	35

Neocate Junior Mixing Instructions

Neocate Junior may be mixed with cool or warm water. Do not boil formula or use microwave to prepare or warm the formula. 1) Measure the required amount of warm or cool water into a container. 2) Add the prescribed amount of Neocate Junior powder to the water. 3) Cover and shake until the powder has dissolved. Prepared formula may be stored in refrigerator and used within 24 hours of mixing. Shake before use. Formula may be kept at room temperature for 4 hours. Once feeding begins, use within 1 hour. May be used for oral or tube feeding.

Storage: Do not store at extreme temperatures. Cover and store opened cans in a cool, dry place (not a refrigerator). Use within one month after opening.

Nutricia North America recommends using a gram scale for greatest accuracy. The scoops Nutricia provides are validated for use with Nutricia formulas.

† Values for volume in fluid ounces are rounded – Milliliter (mL) values are more accurate.

‡ Scoops should be level and unpacked.

To obtain additional information, visit Neocate.com.

Check can label for the most up-to-date mixing instructions.

ZNMP 07/2013



9497350-20023